arXiv:1003.3500v1 [math.FA] 18 Mar 2010

Construction of Symmetric Complex Tight d-wavelet Frames from Pseudo Splines via Matrix Extension with Symmetry

Xiaosheng Zhuang*[,a,1]

[a]*Department of Mathematical and Statistical Sciences, University of Alberta, Edmonton, Alberta, Canada T6G 2G1.*

Abstract

Key words: complex wavelets, pseudo spline, symmetry, framelets, vanishing moments
2000 MSC: 42C40, 41A05, 42C15, 65T60

---

1. Introduction and Motivation

Redundant wavelet systems have been proved to be quit useful in many applications, for examples, signal denoising, image processing, numerical algorithm, and so on. As a redundant system, it can possess many desirable properties such as symmetry, short support, high vanishing moments, and so on, simultaneously. In this paper, we are interested in the construction of tight wavelet frames with such desirable properties. Other than redundancy, computational efficiency is also desired in applications, which means a wavelet system should have as few generators as possible. It is well known that the construction of a wavelet system from a refinable function can be formulated as a matrix extension problem. In order to have fewer generators for a tight wavelet frame, we shall employ the algorithm for matrix extension with symmetry.

Let us first introduce some notation and definitions. We say that d is a *dilation factor* if $\mathsf{d} > 1$ is an integer. Throughout this paper, d is fixed and denotes a dilation factor. We say that $\phi : \mathbb{R} \mapsto \mathbb{C}$ is a d-*refinable function* if

$$\phi = \mathsf{d} \sum_{k\in\mathbb{Z}} a_0(k)\phi(\mathsf{d} \cdot -k), \tag{1.1}$$

where $a_0 : \mathbb{Z} \mapsto \mathbb{C}$ is a finitely supported sequence on $\mathbb{Z}$, called the *low-pass filter* for ϕ. The *symbol* of a_0 is given by $\mathsf{a}_0(z) := \sum_{k\in\mathbb{Z}} a_0(k)z^k$. The *coefficient support* of a_0 is defined by $\operatorname{coeffsupp}(a_0) := [m, n]$, where $a_0(m)a_0(n) \neq 0$ and $a_0(k) = 0$ for all

---

*Corresponding author, `http://www.ualberta.ca/~xzhuang`
Email address: `xzhuang@math.ualberta.ca` (Xiaosheng Zhuang)
[1] Research was supported in part by NSERC Canada under Grant RGP 228051.

$k \notin [m, n]$. In frequency domain, the refinement equation in (1.1) can be rewritten as

$$\hat{\phi}(\mathsf{d}\xi) = \widehat{a_0}(\xi)\hat{\phi}(\xi), \quad \xi \in \mathbb{R}, \tag{1.2}$$

where $\widehat{a_0}$ is the *Fourier series* of a_0 given by

$$\widehat{a_0}(\xi) := \sum_{k\in\mathbb{Z}} a_0(k)e^{-ik\xi} = \mathsf{a}_0(e^{-i\xi}), \quad \xi \in \mathbb{R}. \tag{1.3}$$

The Fourier transform $\hat{f}$ of $f \in L_1(\mathbb{R})$ is defined to be $\hat{f}(\xi) = \int_{\mathbb{R}} f(t)e^{-it\xi}dt$ and can be extended to square integrable functions and tempered distributions. A low-pass filter a_0 is *orthogonal* if $\sum_{j=0}^{\mathsf{d}-1} |\widehat{a_0}(\xi + 2\pi j/\mathsf{d})|^2 = 1$ for all $\xi \in \mathbb{R}$.

Usually, a wavelet system is generated by some wavelet function $\psi^\ell, \ell = 1, \dots, L$ from a d-refinable function ϕ as follows:

$$\widehat{\psi^\ell}(\mathsf{d}\xi) = \widehat{a_\ell}(\xi)\hat{\phi}(\xi), \quad \ell = 1, \dots, L, \tag{1.4}$$

where each $a_\ell : \mathbb{Z} \mapsto \mathbb{C}$ is a finitely supported sequence on $\mathbb{Z}$, called the *high-pass filter* for ψ^ℓ, $\ell = 1, \dots, L$.

We says that $\{\psi^1, \dots, \psi^L\}$ generates a d-*wavelet frame* in $L_2(\mathbb{R})$ if $\{\psi^\ell_{j,k} := \mathsf{d}^{j/2}\psi^\ell(\mathsf{d}^j \cdot -k) : j, k \in \mathbb{Z}, \ell = 1, \dots, L\}$ is a frame in $L_2(\mathbb{R})$, that is, there exists two positive constants C_1, C_2 such that

$$C_1\|f\|^2_{L_2(\mathbb{R})} \leqslant \sum_{\ell=1}^{L} \sum_{j\in\mathbb{Z}} \sum_{k\in\mathbb{Z}} |\langle f, \psi^\ell_{j,k}\rangle|^2 \leqslant C_2\|f\|^2_{L_2(\mathbb{R})} \quad \forall f \in L_2(\mathbb{R}), \tag{1.5}$$

where $|\langle f, \psi^\ell_{j,k}\rangle|^2 = \langle f, \psi^\ell_{j,k}\rangle\langle\psi^\ell_{j,k}, f\rangle$ and $\langle\cdot,\cdot\rangle$ is the inner product in $L_2(\mathbb{R})$ defined by

$$\langle f, g\rangle = \int_{\mathbb{R}} f(t)\overline{g(t)}dt, \quad f, g \in L_2(\mathbb{R}).$$

If $C_1 = C_2 = 1$ in (1.5), we say that $\{\psi^1, \dots, \psi^L\}$ generates a *tight* d-*wavelet frame* in $L_2(\mathbb{R})$.

An important property of a wavelet system is its order of vanishing moments. We say that $\{\psi^1, \dots, \psi^L\}$ has *vanishing moments* of order n if

$$\int_{\mathbb{R}} t^k\psi^\ell(t)dt = 0 \qquad k = 0, \dots, n-1, \forall \ell = 1, \dots, L. \tag{1.6}$$

(1.6) is equivalent to saying that $\frac{d^k}{dt^k}\widehat{\psi^\ell}(0) = 0$ for all $k = 0, \dots, n-1$ and $\ell = 1, \dots, L$.

Let ϕ be a compactly supported d-refinable function in $L_2(\mathbb{R})$ associated with a low-pass filter a_0 such that $\hat{\phi}(0) = 1$. Suppose there exist high-pass filters $a_1, \dots, a_L$ such that

$$\sum_{\ell=0}^{L} \widehat{a_\ell}\overline{\widehat{a_\ell}(\cdot + 2\pi k/\mathsf{d})} = \delta_k, \quad k = 0, \dots, \mathsf{d}-1, \tag{1.7}$$

where δ is the *Dirac sequence* such that $\delta(0) = 1$ and $\delta(k) = 0$ for all $k \neq 0$. Define $\psi^\ell, \ell = 1, \dots, L$ as in (1.4). Then $\{\psi^1, \dots, \psi^L\}$ generates a tight d-wavelet frame, see [11, 10]. Moreover, if the low-pass filter a_0 satisfies

$$1 - |\widehat{a_0}(\xi)|^2 = O(|\xi|^{2n}), \tag{1.8}$$

which means $1-|\widehat{a_0}(\xi)|^2$ has *zero of order* $2n$ near the origin, then $\{\psi^1,\dots,\psi^L\}$ given by (1.4) has vanishing moments of order n, see [4].

Now, we are ready to introduce our main result of this paper on the construction of symmetric complex tight d-wavelet frame. Throughout this paper, we shall denote $P_{m,n}(y)$ a polynomial of degree $n-1$ as follows:

$$P_{m,n}(y)=\sum_{j=0}^{n-1}\left[\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j}\ \prod_{k=1}^{\mathsf{d}-1}\binom{m-1+j_k}{j_k}\sin(k\pi/\mathsf{d})^{-2j_k}\right]y^j. \tag{1.9}$$

Theorem 1. *Let d be a dilation factor. Let $m,n\in\mathbb{N}$ be positive integers such that $2n-1\leqslant m$. Let $P_{m,n}(y)$ be the polynomial defined in* (1.9)*. Then*

$$P_{m,2n-1}(y)>0 \qquad \forall y\in\mathbb{R}. \tag{1.10}$$

Let $z_1,\overline{z_1},\dots,z_{n-1},\overline{z_{n-1}}\in\mathbb{C}\setminus\mathbb{R}$ be all the complex roots of $P_{m,2n-1}$. Then

$$P_{m,2n-1}(y)=|Q_{m,n}(y)|^2, \tag{1.11}$$

where $Q_{m,n}(y)=c(y-z_1)\cdots(y-z_{n-1})$ with $c=(-1)^{n-1}(z_1\cdots z_{n-1})^{-1}$. Define a low-pass filter a_0 by

$$\widehat{a_0}(\xi):=e^{i\lfloor\frac{m(\mathsf{d}-1)}{2}\rfloor\xi}\left(\frac{1+e^{-i\xi}+\cdots+e^{-i(\mathsf{d}-1)\xi}}{\mathsf{d}}\right)^m Q_{m,n}(\sin^2(\xi/2)), \tag{1.12}$$

where $\lfloor\cdot\rfloor$ is the floor operation. Then

$$\widehat{a_0}(-\xi)=e^{i(1-\varepsilon)\xi}\widehat{a_0}(\xi)\quad \textit{with}\quad \varepsilon=m(\mathsf{d}-1)-2\lfloor\frac{m(\mathsf{d}-1)}{2}\rfloor, \tag{1.13}$$

and

$$\operatorname{coeffsupp}(a_0)=[-\lfloor\frac{m(\mathsf{d}-1)}{2}\rfloor-n+1,\lfloor\frac{m(\mathsf{d}-1)}{2}\rfloor+n-1+\varepsilon].$$

Let ϕ be the standard d-refinable function associated with the low-pass filter a_0, that is, $\widehat{\phi}(\xi):=\prod_{j=1}^{\infty}\widehat{a_0}(\mathsf{d}^{-j}\xi)$. Then ϕ is a compactly supported d-refinable function in $L_2(\mathbb{R})$ with symmetry $\phi(\frac{1-\varepsilon}{\mathsf{d}-1}-\cdot)=\phi$. Moreover, one can construct high-pass filters $a_1,\dots,a_L$, $L\in\{\mathsf{d}-1,\mathsf{d},\mathsf{d}+1\}$, with symmetry by Algorithm 2 such that (1.7) *holds. Define $\psi^1,\dots,\psi^L$ as in* (1.4)*. Then $\{\psi^1,\dots,\psi^L\}$ generates a tight d-wavelet frame in $L_2(\mathbb{R})$ and $\{\psi^1,\dots,\psi^L\}$ has vanishing moments of order $2n-1$. Furthermore, $|\operatorname{coeffsupp}(a_\ell)|\leqslant|\operatorname{coeffsupp}(a_0)|$ and $\psi^\ell(\frac{1-\varepsilon}{\mathsf{d}-1}+\eta^\ell-\cdot)=\varepsilon^\ell\psi^\ell$ for some $\eta^\ell\in\{1,0,-1\}$ and $\varepsilon^\ell\in\{-1,1\}$, $\ell=1,\dots,L$. In particular, for $m=2n-1$, $L=\mathsf{d}-1$ and $\{\psi^1,\dots,\psi^{\mathsf{d}-1}\}$ generates an orthonormal wavelet basis with symmetry for $L_2(\mathbb{R})$.*

Theorem 1 is summarized from the following sections. In section 2, we shall inspect the important properties of $P_{m,n}(y)$ and show that for any dilation factor d and $m,n\in\mathbb{N}$, we can construct a symmetric complex d-refinable function from $P_{m,2n-1}(y)$, called ***complex*** *d-refinable pseudo spline of type I with order* $(m,2n-1)$. Moreover, we prove that the shifts of all such complex pseudo splines are linearly independent.

In section 3, we introduce the general problem of matrix extension with symmetry and present a step-by-step algorithm for the matrix extension with symmetry. In section 4, we shall apply our matrix extension algorithm and show that we can construct symmetric complex tight d-wavelet frames from symmetric complex pseudo spline with only d wavelet generators. Moreover, we shall provide several examples to demonstrate our results and algorithms. Conclusions will be given in last section.

2. Symmetric Complex Refinable Functions from Pseudo Splines

In this section, we shall restrict our attention to the construction of d-refinable functions from which wavelet system with high vanishing moments can be derived. In other words, we need to design low-pass filters a such that a satisfies (1.8) for some $n \in \mathbb{N}$.

To guarantee that the d-refinable function ϕ associated with a has certain regularity, usually the low-pass filter a satisfies the *sum rules of order m* for some $m \in \mathbb{N}$:

$$(1 + e^{-i\xi} + \cdots + e^{-i(\mathsf{d}-1)\xi})^m \mid \hat{a}(\xi). \tag{2.1}$$

That is, $\widehat{a}$ is of the form:

$$\hat{a}(\xi) = e^{i\lfloor \frac{m(\mathsf{d}-1)}{2} \rfloor} \left(\frac{1 + e^{-i\xi} + \cdots + e^{-i(\mathsf{d}-1)\xi}}{\mathsf{d}} \right)^m \widehat{L}(\xi), \tag{2.2}$$

for some 2π-periodic trigonometric polynomial $\widehat{L}$ with $\widehat{L}(0) = 1$. For $\widehat{L}(\xi) \equiv 1$. $\widehat{a}(\xi)$ is the low-pass filter for B-spline of order m: $\widehat{B_m}(\xi) = e^{i\lfloor m/2 \rfloor \xi}(1 - e^{-i\xi})^m/(i\xi)^m$.

Define a function h by

$$h(y) := \prod_{k=1}^{\mathsf{d}-1} \left(1 - \frac{y}{\sin^2(k\pi/\mathsf{d})} \right), \quad y \in \mathbb{R}. \tag{2.3}$$

We have

$$h(\sin^2(\xi/2)) = \frac{|1 + \cdots + e^{-i(\mathsf{d}-1)\xi}|^2}{\mathsf{d}^2} = \frac{\sin^2(\mathsf{d}\xi/2)}{\mathsf{d}^2 \sin^2(\xi/2)} \tag{2.4}$$

and

$$h(y)^{-m} = \left[\prod_{k=1}^{\mathsf{d}-1} \left(\sum_{j_k=0}^{\infty} \frac{y^{j_k}}{\sin^{2j_k}(k\pi/\mathsf{d})} \right) \right]^{-m} = \sum_{j=0}^{\infty} c_{m,j} y^j, \quad |y| < \sin^2(\pi/\mathsf{d}), \tag{2.5}$$

where

$$c_{m,j} = \sum_{j_1+\cdots+j_{\mathsf{d}-1}=j} \prod_{k=1}^{\mathsf{d}-1} \binom{m-1+j_k}{j_k} \sin(k\pi/\mathsf{d})^{-2j_k}, \quad j \in \mathbb{N}. \tag{2.6}$$

Note that $P_{m,n}(y) = \sum_{j=0}^{n-1} c_{m,j} y^j$. Consequently, we have the following result.

Lemma 2. *Let $m, n \in \mathbb{N}$ be such that $n \leqslant m$; let $P_{m,n}$ and h be polynomials defined as in* (1.9) *and* (2.3)*, respectively. Then $P_{m,n}(\sin^2(\xi/2))$ is the unique positive trigonometric polynomial of minimal degree such that*

$$1 - h(\sin^2(\xi/2))^m P_{m,n}(\sin^2(\xi/2)) = O(|\xi|^{2n}). \tag{2.7}$$

Proof. Suppose there exist trigonometric polynomial $\hat{g}(\xi)$ such that

$$1 - h(\sin^2(\xi/2))^m g(\xi) = O(|\xi|^{2n}).$$

Observing that the first $(n-1)th$-degree Taylor polynomial of $h(y)^{-m}$ at $y = 0$ is $P_{m,n}(y)$, we have

$$g(\xi) = h(\sin^2(\xi/2))^{-m}[1 + O(|\xi|^{2n})] = P_{m,n}(\sin^2(\xi/2)) + O(|\sin^2(\xi/2)|^n).$$

Moreover, it is easily seen that the coefficients of $P_{m,n}(y)$ are all positive. Consequently, $P_{m,n}(\sin^2(\xi/2))$ is the unique positive trigonometric polynomial of minimal degree such that (2.7) holds. ■

For $m, n \in \mathbb{N}$ such that $n \leqslant m$, let $\widehat{{}_{II}a}(\xi) := h(\sin^2(\xi/2)^m P_{m,n}(\sin^2(\xi/2))$. Then the d-refinable function ${}_{II}\phi$ associated with ${}_{II}a$ by (1.1) is called the *pseudo spline of type II*. By Lemma 2, using Riesz Lemma, one can derive a low-pass filter ${}_{I}a$ from ${}_{II}a$ such that $|\widehat{{}_{I}a}(\xi)|^2 = \widehat{{}_{II}a}(\xi)$. The d-refinable function ${}_{I}\phi$ associated with such ${}_{I}a$ by (1.1) is referred as *real pseudo spline of type I*. Interesting readers can refer to [**? ? ? ?**] for more details on this subject for the special case $\mathsf{d} = 2$.

Note that ${}_{I}a$ satisfies (1.8). One can construct high-pass filters $b^1, \dots, b^L$ from ${}_{I}a$ such that (1.7) holds. Then $\psi^1, \dots, \psi^L$ defined by (1.4) are real-valued functions. $\{\psi^1, \dots, \psi^L\}$ has vanishing moment of order n and generates a tight d-wavelet frame. However, $\{\psi^1, \dots, \psi^L\}$ is not necessary symmetric since the low-pass filter ${}_{I}a$ from ${}_{II}a$ via Riesz lemma might not posses any symmetry pattern. In what follows, we shall consider complex-valued wavelet generators and show that we can achieve symmetry for any odd integer $n \in N$.

The next lemma is needed later, which generalizes [**?** , Lemma 2.1].

Lemma 3. *Let $P_{m,n}(y)$ be defined as in* (1.9)*. Then*

$$P'_{m,n}(y) = m \sum_{\ell=1}^{\mathsf{d}-1} \frac{1}{(1 - \frac{y}{\sin^2(\frac{\ell\pi}{\mathsf{d}})})} \left[\frac{P_{m,n}(y)}{\sin^2(\frac{\ell\pi}{\mathsf{d}})} - c^{\ell}_{n-1} y^{n-1} \right], \tag{2.8}$$

where

$$c^{\ell}_{n-1} = \sum_{j_1+\cdots+j_{\mathsf{d}-1}=n-1} \left[\binom{m+j_\ell}{j_\ell} \sin(\frac{\ell\pi}{\mathsf{d}})^{-2(j_\ell+1)} \prod_{k=1,k\neq\ell}^{\mathsf{d}-1} \binom{m-1+j_k}{j_k} \sin(\frac{k\pi}{\mathsf{d}})^{-2j_k} \right].$$

Proof. By (1.9), we have

$$\begin{aligned}
P'_{m,n}(y) &= \sum_{j=0}^{n-2} \left[\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j+1} \prod_{k=1}^{\mathsf{d}-1} \binom{m-1+j_k}{j_k} \sin(\frac{k\pi}{\mathsf{d}})^{-2j_k} \right] (j+1) y^j \\
&= m \sum_{j=0}^{n-2} \left[\sum_{\ell=1}^{\mathsf{d}-1} \sum_{j_1+\cdots+j_{\mathsf{d}-1}=j+1} \frac{j_\ell}{m} \prod_{k=1}^{\mathsf{d}-1} \binom{m-1+j_k}{j_k} \sin(\frac{k\pi}{\mathsf{d}})^{-2j_k} \right] y^j \\
&= m \sum_{\ell=1}^{\mathsf{d}-1} \sum_{j=0}^{n-2} \left[\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j} \binom{m+j_\ell}{j_\ell} \sin(\frac{\ell\pi}{\mathsf{d}})^{-2(j_\ell+1)} \prod_{k=1,k\neq\ell}^{\mathsf{d}-1} \binom{m-1+j_k}{j_k} \sin(\frac{k\pi}{\mathsf{d}})^{-2j_k} \right] y^j \\
&=: m \sum_{\ell=1}^{\mathsf{d}-1} Q_\ell(y),
\end{aligned}$$

where

$$Q_\ell(y) := \sum_{j=0}^{n-2}\left[\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j}\binom{m+j_\ell}{j_\ell}\sin(\frac{\ell\pi}{\mathsf{d}})^{-2(j_\ell+1)}\prod_{k=1,k\neq\ell}^{\mathsf{d}-1}\binom{m-1+j_k}{j_k}\sin(\frac{k\pi}{\mathsf{d}})^{-2j_k}\right]y^j$$

Note that

$$\begin{aligned}
&\frac{y}{\sin^2(\frac{\ell\pi}{\mathsf{d}})}Q_\ell(y)\\
&=\sum_{j=0}^{n-2}\left[\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j}\binom{m+j_\ell}{j_\ell}\sin(\frac{\ell\pi}{\mathsf{d}})^{-2(j_\ell+2)}\prod_{k=1,k\neq\ell}^{\mathsf{d}-1}\binom{m-1+j_k}{j_k}\sin(\frac{k\pi}{\mathsf{d}})^{-2j_k}\right]y^{j+1}\\
&=\sum_{j=1}^{n-1}\left[\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j}\binom{m-1+j_\ell}{j_\ell-1}\sin(\frac{\ell\pi}{\mathsf{d}})^{-2(j_\ell+1)}\prod_{k=1,k\neq\ell}^{\mathsf{d}-1}\binom{m-1+j_k}{j_k}\sin(\frac{k\pi}{\mathsf{d}})^{-2j_k}\right]y^j.
\end{aligned}$$

Using $\binom{m+j_\ell}{j_\ell}-\binom{m-1+j_\ell}{j_\ell-1}=\binom{m-1+j_\ell}{j_\ell}$, it is easy to deduce that

$$\left(1-\frac{y}{\sin^2(\frac{\ell\pi}{\mathsf{d}})}\right)Q_\ell(y)=\frac{P_{m,n}(y)}{\sin^2(\frac{\ell\pi}{\mathsf{d}})}-c_{n-1}^{\ell}y^{n-1},$$

where

$$c_{n-1}^{\ell}=\sum_{j_1+\cdots+j_{\mathsf{d}-1}=n-1}\left[\binom{m+j_\ell}{j_\ell}\sin(\frac{\ell\pi}{\mathsf{d}})^{-2(j_\ell+1)}\prod_{k=1,k\neq\ell}^{\mathsf{d}-1}\binom{m-1+j_k}{j_k}\sin(\frac{k\pi}{\mathsf{d}})^{-2j_k}\right].$$

Consequently, (2.8) holds. ■

By the above lemma, we have the following result regarding the positiveness of $P_{m,n}(y)$, which generalizes [**?** , Theorem 5] and [**?** , Theorem 2.4] and proves (1.10).

Theorem 4. *Let $m, n \in \mathbb{N}$ be such that $n \leqslant m$. Then $P_{m,n}(y) > 0$ for all $y \in \mathbb{R}$ if and only if n is an odd number.*

Proof. It is easily seen that $P_{m,n}(y) > 0$ for all $y \geqslant 0$ due to the positiveness of its coefficients and $\lim_{y\to-\infty} P_{m,n}(y) = -\infty$ when n is even. Hence n must be an odd number if $P_{m,n}(y) > 0$ for all $y \in \mathbb{R}$.

On the other hand, suppose n is an odd number. Let $y_0 < 0$ be any stationary point of $P_{m,n}(y)$. Then by Lemma 3, we have

$$0 = P'_{m,n}(y_0) = m\sum_{\ell=1}^{\mathsf{d}-1}\frac{1}{(1-\frac{y_0}{\sin^2(\frac{\ell\pi}{\mathsf{d}})})}\left[\frac{P_{m,n}(y_0)}{\sin^2(\frac{\ell\pi}{\mathsf{d}})}-c_{n-1}^{\ell}y_0^{n-1}\right],$$

which implies that $P_{m,n}(y_0) > 0$. Consequently, $P_{m,n}(y) > 0$ for all $y < 0$. This completes our proof. ■

Now, by $P_{m,2n-1}(y) > 0$ for all $y \in \mathbb{R}$ and $2n-1 \leqslant m$, $P_{m,2n-1}(y)$ can only have complex roots. Hence, we must have

$$P_{m,2n-1}(y) = c_0 \prod_{j=1}^{n-1} (y - z_j)(y - \overline{z_j}), \quad z_1, \overline{z_1}, \ldots, z_{n-1}, \overline{z_{n-1}} \in \mathbb{C} \setminus \mathbb{R}.$$

Define $Q_{m,n}(y) = c \prod_{j=1}^{n-1}(y - z_j)$ as in (1.11) and $\widehat{a_0}(\xi)$ as in (1.12). Then it is easy to check that the symmetry pattern of a_0 satisfies (1.13). We shall refer the d-refinable function $\phi_{m,n}$ associated with the low-pass filter a_0 as *complex pseudo spline of type I.*

Now, we have the following result which shall play an important role in our construction of tight d-wavelet frame in section 3.

Corollary 5. *Let $m, n \in \mathbb{N}$ be such that $2n-1 \leqslant m$ and a_0 be defined as in* (1.12)*. Then*

$$1 - \sum_{j=0}^{\mathsf{d}-1} |\widehat{a_0}(\xi + 2\pi j/\mathsf{d})|^2 = |\widehat{b_0}(\mathsf{d}\xi)|^2, \tag{2.9}$$

for some 2π-periodic trigonometric function $\widehat{b_0}(\xi)$ with real coefficients. In particular, $\widehat{b_0^{2n-1,n}}(\xi) \equiv 0$ and $|\widehat{b_0^{2n,n}}(\xi)|^2 = c_{2n,2n-1}[\sin^2(\xi/2)/\mathsf{d}^2]^{2n-1}$ with $c_{2n,2n-1}$ being the coefficient given in (2.6).

Proof. We first show the $1 - \sum_{j=0}^{\mathsf{d}-1} |\widehat{a_0}(\xi + 2\pi j/\mathsf{d})|^2 \geqslant 0$ for all $\xi \in \mathbb{R}$. Let $y_j := \sin^2(\xi/2 + \pi j/\mathsf{d})$, $j = 0, \ldots, \mathsf{d}-1$. Since $|\widehat{a_0}(\xi)|^2 = h(\sin^2(\xi/2))^m P_{m,2n-1}(\sin^2(\xi/2))$, we have

$$\begin{aligned}
1 - \sum_{j=0}^{\mathsf{d}-1} |\widehat{a_0}(\xi + 2\pi j/\mathsf{d})|^2 &= 1 - \sum_{j=0}^{\mathsf{d}-1} h(y_j)^m P_{m,2n-1}(y_j) \\
&= 1 - \sum_{j=0}^{\mathsf{d}-1} h(y_j)^m P_{m,m}(y_j) + \sum_{j=0}^{\mathsf{d}-1} h(y_j)^m \sum_{k=2n-1}^{m-1} c_{m,k} y_j^k \\
&= \sum_{j=0}^{\mathsf{d}-1} h(y_j)^m \sum_{k=2n-1}^{m-1} c_{m,k} y_j^k \geqslant 0.
\end{aligned}$$

The last equality follows from the fact that the low-pass filter a satisfying $|\hat{a}(\xi)|^2 = h(y_0)^m P_{m,m}(y_0)$ is an orthogonal low-pass filter , which satisfies $\sum_{j=0}^{\mathsf{d}-1} h(y_j)^m P_{m,m}(y_j) = 1$ ([**?**]). Now, by that $1 - \sum_{j=0}^{\mathsf{d}-1} |\widehat{a_0}(\xi + 2\pi j/\mathsf{d})|^2 \geqslant 0$ is of period $2\pi/\mathsf{d}$, (2.9) follows from Riesz Lemma.

Obviously, $\widehat{b_0^{2n-1,n}}(\xi) \equiv 0$ since $a_0^{2n-1,n}$ is an orthogonal low-pass filter. And from above, noting that $h(y_j)y_j = \sin^2(\mathsf{d}\xi/2)/\mathsf{d}^2$. we have

$$\begin{aligned}
|\widehat{b_0^{2n,2n-1}}(\mathsf{d}\xi)|^2 &= c_{2n,2n-1} \sum_{j=0}^{\mathsf{d}-1} h(y_j)^{2n} y_j^{2n-1} = c_{2n,2n-1} \sum_{j=0}^{\mathsf{d}-1} (h(y_j)y_j)^{2n-1}) h(y_j) \\
&= c_{2n,2n-1}[\sin^2(\mathsf{d}\xi/2)/\mathsf{d}^2]^{2n-1} \sum_{j=0}^{\mathsf{d}-1} h(y_j) P_{1,1}(y_j) \\
&= c_{2n,2n-1}[\sin^2(\mathsf{d}\xi/2)/\mathsf{d}^2]^{2n-1}.
\end{aligned}$$

■

Next, we shall discuss the linear independence of the pseudo spline of type I (real and complex) and type II. Linear independence is an important issue in both approximation theory and wavelet analysis. The linear independence of the integer shifts of a refinable function is a necessary and sufficient condition for the existence of a compactly supported dual refinable function, see [**?**]. For a compactly supported function $\phi \in L_2(\mathbb{R})$, we say that the shifts of ϕ are *linearly independent* if

$$\sum_{j\in\mathbb{Z}} c(j)\phi(\cdot - j) = 0 \quad \text{implies} \quad c(j) = 0\,\forall j \in \mathbb{Z}. \tag{2.10}$$

We say that the shifts of ϕ are *stable* if there exists two positive constants C_1 and C_2 such that

$$C_1 \sum_{j\in\mathbb{Z}} |c(j)|^2 \leqslant \|\sum_{j\in\mathbb{Z}} c(j)\phi(\cdot - j)\|^2_{L_2(\mathbb{R})} \leqslant C_2 \sum_{j\in\mathbb{Z}} |c(j)|^2, \tag{2.11}$$

for all finitely supported sequences $c : \mathbb{Z} \mapsto \mathbb{C}$.

It is known in [**?**] that the shifts of a compacly supported function $\phi \in L_2(\mathbb{R})$ is linearly independent (or stable) if and only if

$$\text{span}\{\hat{\phi}(\xi + 2\pi k) \ : \ k \in \mathbb{Z}\} \neq 0 \ \text{ for all } \ \xi \in \mathbb{C} \,(\text{or for all } \xi \in \mathbb{R}). \tag{2.12}$$

Here, the *Fourier–Lapalce transform* of ϕ is defined to be

$$\hat{\phi}(\xi) = \int_{\mathbb{R}} \phi(t) e^{-i\xi t} dt, \quad \xi \in \mathbb{C}. \tag{2.13}$$

Consequently, if the shifts of a compactly supported function $\phi \in L_2(\mathbb{R})$ are linearly independent, then the shifts of ϕ must be stable in $L_2(\mathbb{R})$.

The following lemma is needed later to prove the linear independence of the shifts of pseudo spline of type I or type II.

Lemma 6. *Let $m, n \in \mathbb{N}$ be such that $1 < n < m$. Let $c_{m,j}$ be the coefficient of $P_{m,n}(y)$ defined in* (2.6). *Then*

$$2c_{m,j-1} < c_{m,j}, \quad j = 1, \ldots n - 1. \tag{2.14}$$

Proof. First, it is easy to show that $2\binom{m-1+j}{j} < \binom{m+j}{j+1}$ for $j = 0, \ldots, n-2$. Then

$$
\begin{aligned}
2c_{m,j-1} &= 2\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j-1}\prod_{k=1}^{\mathsf{d}-1}\binom{m-1+j_k}{j_k}\sin(\frac{k\pi}{\mathsf{d}})^{-2j_k} \\
&= \frac{1}{\mathsf{d}-1}\sum_{\ell=1}^{\mathsf{d}-1}\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j-1} 2\binom{m-1+j_\ell}{j_\ell}\sin(\frac{\ell\pi}{\mathsf{d}})^{-2j_\ell}\prod_{k=1,k\neq\ell}^{\mathsf{d}-1}\binom{m-1+j_k}{j_k}\sin(\frac{k\pi}{\mathsf{d}})^{-2j_k} \\
&< \frac{1}{\mathsf{d}-1}\sum_{\ell=1}^{\mathsf{d}-1}\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j-1} \binom{m+j_\ell}{j_\ell+1}\sin(\frac{\ell\pi}{\mathsf{d}})^{-2(j_\ell+1)}\prod_{k=1,k\neq\ell}^{\mathsf{d}-1}\binom{m-1+j_k}{j_k}\sin(\frac{k\pi}{\mathsf{d}})^{-2j_k} \\
&= \frac{1}{\mathsf{d}-1}\sum_{\ell=1}^{\mathsf{d}-1}\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j} \binom{m-1+j_\ell}{j_\ell}\sin(\frac{\ell\pi}{\mathsf{d}})^{-2(j_\ell)}\prod_{k=1,k\neq\ell}^{\mathsf{d}-1}\binom{m-1+j_k}{j_k}\sin(\frac{k\pi}{\mathsf{d}})^{-2j_k} \\
&= \frac{1}{\mathsf{d}-1}\sum_{\ell=1}^{\mathsf{d}-1}\sum_{j_1+\cdots+j_{\mathsf{d}-1}=j}\prod_{k=1}^{\mathsf{d}-1}\binom{m-1+j_k}{j_k}\sin(\frac{k\pi}{\mathsf{d}})^{-2j_k} \\
&= c_{m,j}.
\end{aligned}
$$

■

Due to $\widehat{{}_{II}\phi}(\xi) = \widehat{{}_{I}\phi}(\xi)\cdot\widehat{\overline{{}_{I}\phi}}(-\xi)$ for all $\xi \in \mathbb{C}$, the shifts of the pseudo spline of type II are linearly independent will implies the shifts of the pseudo spline of type I are linearly independent as well [**?** , Proposition 1.1]. Moreover, we have the following lemma, whose proof is also similar to [**?** , Lemma 2.1]

Lemma 7. *Let $\phi \in L_2(\mathbb{R})$ be a compactly supported refinable function associated with a low-pass filter a by* (1.1)*. Then, the shifts of ϕ are linearly independent if and only if*

(1) *the shifts of ϕ are stable;*

(2) *the symbol a of a satisfies:* $\{\mathsf{a}(ze^{i2\pi j/\mathsf{d}}) : j = 0, \ldots, \mathsf{d}-1\} \neq 0$ *for all $z \in \mathbb{C}$.*

For a polynomial $P(z) = c_0 + c_1 z + \cdots + c_n z^n$ with real coefficients satisfying $c_n > c_{n-1} > \cdots > c_0 > 0$, it was shown in [**?**] that all zeros of $P(z)$ are containing inside the open unit disk $\{z \in \mathbb{C} : |z| < 1\}$.

Using the above facts, we can prove the following result.

Theorem 8. *The shifts of the pseudo spline of type I (real or complex) or type II are linearly independent.*

Proof. We only need to show that the shifts of the pseudo spline of type II are linearly independent.

Fix $m, n \in \mathbb{N}$ with $1 \leqslant n \leqslant m$. Let the low-pass filter a be given as follows:

$$\hat{a}(\xi) = h(\sin^2(\xi/2))^m P_{m,n}(\sin^2(\xi/2)).$$

Let ϕ be the pseudo spline of type II associated with mask a by (1.1).

For $n = m$, ϕ associated with a is an interpolatory d-refinable function [**?**], i.e., $\phi(j) = \delta(j)$, $j \in \mathbb{Z}$ and $\hat{\phi}(\mathsf{d}\xi) = \hat{a}(\xi)\hat{\phi}(\xi)$. In this case, the shifts of ϕ must be linearly independent.

For $n = 1$, by that the shifts of the B-spline of order $2m$ are stable and the symbol of a is $\mathsf{a}(z) = |1 + \cdots + z^{\mathsf{d}-1}|^{2m}/\mathsf{d}^{2m}$, the fact that the shifts of ϕ associated with a are linearly independent follows by Lemma 7.

Hence, without loss of generality, we can assume that $1 < n < m$. Since $\hat{a}(\xi) \geqslant h(\sin^2(\xi/2))^m$ and the shifts of B-spine of order $2m$ are stable, by (2.12), the shifts of ϕ are stable. We need to prove that the symbol of a satisfies item (2) of Lemma 7. By $\sin^2(\xi/2) = -\frac{(1-z)^2}{4z}$ with $z = e^{-i\xi}$, we have

$$\begin{aligned}
\mathsf{a}(z) &= \left|\frac{1 + z + \cdots + z^{\mathsf{d}-1}}{\mathsf{d}}\right|^{2m} P_{m,n}\left(-\frac{(1-z)^2}{4z}\right) \\
&= \left|\frac{1 + z + \cdots + z^{\mathsf{d}-1}}{\mathsf{d}}\right|^{2m} \sum_{j=0}^{n-1} c_{m,j}\left[-\frac{(1-z)^2}{4z}\right]^j \\
&= \left|\frac{1 + z + \cdots + z^{\mathsf{d}-1}}{\mathsf{d}}\right|^{2m} \sum_{j=0}^{n-1} 2^{-j} c_{m,j}\left[-\frac{(1-z)^2}{2z}\right]^j .
\end{aligned}$$

Now, item (2) of Lemma 7 is equivalent to saying that there is no $z \in \mathbb{C}$ such that $P(ze^{i2\pi j/\mathsf{d}}) = 0$ for all $j = 0, \ldots, \mathsf{d} - 1$, where $P(z) = \sum_{j=0}^{n-1} 2^{-j} c_{m,j}\left[-\frac{(1-z)^2}{2z}\right]^j$. Suppose not, let $z_0 = \rho e^{i\theta} \in \mathbb{C}$ be such that $P(z_j) = 0$ for all $j = 0, \ldots, \mathsf{d} - 1$, where $z_j = \rho e^{i\theta_j}$ with $\theta_j = \theta + 2\pi j/\mathsf{d}$ for $j = 0, \ldots, \mathsf{d} - 1$. Note that

$$\left|-\frac{(1-z_j)^2}{2z_j}\right| = \frac{|1-z_j|^2}{2\rho} = \frac{\rho^2+1}{2\rho} - \cos(\theta_j) \geqslant 1 - \cos(\theta_j), \quad j = 0, \ldots, \mathsf{d} - 1.$$

Consequently, $\max_{0\leqslant j\leqslant \mathsf{d}-1}\left|-\frac{(1-z_j)^2}{2z_j}\right| \geqslant 1$. This contradicts to the fact that all roots of the polynomial $\sum_{j=0}^{n-1} 2^{-j} c_{m,j} z^j$ are containing inside the unit disk $\{z \in \mathbb{C} : |z| < 1\}$ due to its coefficients $2^j c_{m,j}$, $j = 0, \ldots, n - 1$ satisfying $2^{-j+1} c_{m,j-1} < 2^{-j} c_{m,j}$ for all $j = 1, \ldots, n - 1$. We are done. ■

3. Matrix Extension with Symmetry

In this section, we shall introduce the matrix extension problem, which plays an important role in our construction of tight d-wavelet frame. In order to state the matrix extension problem and our main results, let us introduce some notation and definitions first.

Let $\mathsf{p}(z) = \sum_{k\in\mathbb{Z}} p_k z^k, z \in \mathbb{C}\backslash\{0\}$ be a Laurent polynomial with complex coefficients $p_k \in \mathbb{C}$ for all $k \in \mathbb{Z}$. We say that p has *symmetry* if its coefficient sequence $\{p_k\}_{k\in\mathbb{Z}}$ has symmetry; more precisely, there exist $\varepsilon \in \{-1, 1\}$ and $c \in \mathbb{Z}$ such that

$$p_{c-k} = \varepsilon p_k, \qquad \forall\, k \in \mathbb{Z}. \tag{3.1}$$

If $\varepsilon = 1$, then p is symmetric about the point $c/2$; if $\varepsilon = -1$, then p is antisymmetric about the point $c/2$. Symmetry of a Laurent polynomial can be conveniently expressed using a symmetry operator $\mathcal{S}$ defined by

$$\mathcal{S}\mathsf{p}(z) := \frac{\mathsf{p}(z)}{\mathsf{p}(1/z)}, \qquad z \in \mathbb{C}\backslash\{0\}. \tag{3.2}$$

When p is not identically zero, it is evident that (3.1) holds if and only if $\mathcal{S}\mathsf{p}(z) = \varepsilon z^c$. For the zero polynomial, it is very natural that $\mathcal{S}0$ can be assigned any symmetry pattern; that is, for every occurrence of $\mathcal{S}0$ appearing in an identity in this paper, $\mathcal{S}0$ is understood to take an appropriate choice of εz^c for some $\varepsilon \in \{-1, 1\}$ and $c \in \mathbb{Z}$ so that the identity holds. If P is an $r \times s$ matrix of Laurent polynomials with symmetry, then we can apply the operator $\mathcal{S}$ to each entry of P, that is, $\mathcal{S}\mathsf{P}$ is an $r \times s$ matrix such that $[\mathcal{S}\mathsf{P}]_{j,k} := \mathcal{S}([\mathsf{P}]_{j,k})$, where $[\mathsf{P}]_{j,k}$ denotes the (j, k)-entry of the matrix P throughout the chapter.

For two matrices P and Q of Laurent polynomials with symmetry, even though all the entries in P and Q have symmetry, their sum $\mathsf{P} + \mathsf{Q}$, difference $\mathsf{P} - \mathsf{Q}$, or product PQ, if well defined, generally may not have symmetry any more. This is one of the difficulties for matrix extension with symmetry. In order for $\mathsf{P} \pm \mathsf{Q}$ or PQ to possess some symmetry, the symmetry patterns of P and Q should be compatible. For example, if $\mathcal{S}\mathsf{P} = \mathcal{S}\mathsf{Q}$, that is, both P and Q have the same symmetry pattern, then indeed $\mathsf{P} \pm \mathsf{Q}$ has symmetry and $\mathcal{S}(\mathsf{P} \pm \mathsf{Q}) = \mathcal{S}\mathsf{P} = \mathcal{S}\mathsf{Q}$. In the following, we discuss the compatibility of symmetry patterns of matrices of Laurent polynomials. For an $r \times s$ matrix $\mathsf{P}(z) = \sum_{k\in\mathbb{Z}} P_k z^k$, we denote

$$\mathsf{P}^*(z) := \sum_{k\in\mathbb{Z}} P_k^* z^{-k} \quad \text{with} \quad P_k^* := \overline{P_k}^T, \qquad k \in \mathbb{Z}, \tag{3.3}$$

where $\overline{P_k}^T$ denotes the transpose of the complex conjugate of the constant matrix P_k in $\mathbb{C}$. We say that *the symmetry of* P *is compatible* or P *has compatible symmetry*, if

$$\mathcal{S}\mathsf{P}(z) = (\mathcal{S}\theta_1)^*(z)\mathcal{S}\theta_2(z), \tag{3.4}$$

for some $1 \times r$ and $1 \times s$ row vectors θ_1 and θ_2 of Laurent polynomials with symmetry. For an $r \times s$ matrix P and an $s \times t$ matrix Q of Laurent polynomials, we say that (P, Q) *has mutually compatible symmetry* if

$$\mathcal{S}\mathsf{P}(z) = (\mathcal{S}\theta_1)^*(z)\mathcal{S}\theta(z) \quad \text{and} \quad \mathcal{S}\mathsf{Q}(z) = (\mathcal{S}\theta)^*(z)\mathcal{S}\theta_2(z) \tag{3.5}$$

for some $1 \times r$, $1 \times s$, $1 \times t$ row vectors $\theta_1, \theta, \theta_2$ of Laurent polynomials with symmetry. If (P, Q) has mutually compatible symmetry as in (3.5), then it is easy to verify that their product PQ has compatible symmetry and in fact $\mathcal{S}(\mathsf{PQ}) = (\mathcal{S}\theta_1)^*\mathcal{S}\theta_2$.

For a matrix of Laurent polynomials, another important property is the support of its coefficient sequence. For $\mathsf{P} = \sum_{k\in\mathbb{Z}} P_k z^k$ such that $P_k = \mathbf{0}$ for all $k \in \mathbb{Z}\backslash[m, n]$ with $P_m \neq \mathbf{0}$ and $P_n \neq \mathbf{0}$, we define its coefficient support to be $\mathrm{coeffsupp}(\mathsf{P}) := [m, n]$ and the length of its coefficient support to be $|\mathrm{coeffsupp}(\mathsf{P})| := n - m$. In particular, we define $\mathrm{coeffsupp}(\mathbf{0}) := \emptyset$, the empty set, and $|\mathrm{coeffsupp}(\mathbf{0})| := -\infty$. Also, we use

$\operatorname{coeff}(\mathsf{P}, k) := P_k$ to denote the coefficient matrix (vector) P_k of z^k in P. In this chapter, $\mathbf{0}$ always denotes a general zero matrix whose size can be determined in the context.

The Laurent polynomials that we shall consider in this paper have their coefficients in a subfield $\mathbb{F}$ of the complex field $\mathbb{C}$. Let $\mathbb{F}$ denote a subfield of $\mathbb{C}$ such that $\mathbb{F}$ is closed under the operations of complex conjugate of $\mathbb{F}$ and square roots of positive numbers in $\mathbb{F}$. In other words, the subfield $\mathbb{F}$ of $\mathbb{C}$ satisfies the following properties:

$$\bar{x} \in \mathbb{F} \quad \text{and} \quad \sqrt{y} \in \mathbb{F}, \qquad \forall\, x, y \in \mathbb{F} \quad \text{with} \quad y > 0. \tag{3.6}$$

Two particular examples of such subfields $\mathbb{F}$ are $\mathbb{F} = \mathbb{R}$ (the field of real numbers) and $\mathbb{F} = \mathbb{C}$ (the field of complex numbers). A nontrivial example other than $\mathbb{R}$ and $\mathbb{C}$ that satisfies (3.6) is the field of all algebraic number, i.e., the algebraic closure $\overline{\mathbb{Q}}$ of the rational number $\mathbb{Q}$. A subfield of $\mathbb{R}$ given by $\overline{\mathbb{Q}} \cap \mathbb{R}$ also satisfies (3.6).

Now, we introduce the general matrix extension problem with symmetry. Let r and s denote two positive integers such that $1 \leqslant r \leqslant s$. Let P be an $r \times s$ matrix of Laurent polynomials with coefficients in $\mathbb{F}$ such that $\mathsf{P}(z)\mathsf{P}^*(z) = I_r$ for all $z \in \mathbb{C}\backslash\{0\}$ and the symmetry of P is compatible, where I_r denotes the $r \times r$ identity matrix. The matrix extension problem with symmetry is to find an $s \times s$ square matrix P_e of Laurent polynomials with coefficients in $\mathbb{F}$ and with symmetry such that $[I_r, \mathbf{0}]\mathsf{P}_e = \mathsf{P}$ (that is, the submatrix of the first r rows of P_e is the given matrix P), the symmetry of P_e is compatible, and $\mathsf{P}_e(z)\mathsf{P}_e^*(z) = I_s$ for all $z \in \mathbb{C}\backslash\{0\}$ (that is, P_e is paraunitary). Moreover, in many applications, it is often highly desirable that the coefficient support of P_e can be controlled by that of P in some way.

In the context of wavelet analysis, matrix extension without symmetry has been discussed by Lawton, Lee and Shen in their interesting paper [8] and a simple algorithm has been proposed there to derive a desired matrix P_e from a given row vector P of Laurent polynomials without symmetry. In electronic engineering, an algorithm using the cascade structure for matrix extension without symmetry has been given in [12] for filter banks with perfect reconstruction property. The algorithms in [8, 12] mainly deal with the special case that P is a row vector without symmetry and the coefficient support of the derived matrix P_e indeed can be controlled by that of P. For $\mathbb{F} = \mathbb{R}$ and $r = 1$, matrix extension with symmetry has been considered in [9].

We study the general matrix extension problem with symmetry and have the following results.

Theorem 9. *Let* P *be an* $r \times s$ *matrix of Laurent polynomials with coefficients in a subfield* $\mathbb{F}$ *of* $\mathbb{C}$ *such that* (3.6) *holds. Then* $\mathsf{P}(z)\mathsf{P}^*(z) = I_r$ *for all* $z \in \mathbb{C}\backslash\{0\}$ *and the symmetry of* P *is compatible as in* (3.4)*, if and only if, there exist* $s \times s$ *matrices* $\mathsf{P}_0, \ldots, \mathsf{P}_{J+1}$ *of Laurent polynomials with coefficients in* $\mathbb{F}$ *such that*

(1) $\mathsf{P}_e := \mathsf{P}_{J+1}(z)\mathsf{P}_J(z)\cdots\mathsf{P}_1(z)\mathsf{P}_0(z)$ *is paraunitary:* $\mathsf{P}_e(z)\mathsf{P}_e^*(z) = I_s$*;*

(2) P *can be represented as a product of* $\mathsf{P}_0, \ldots, \mathsf{P}_{J+1}$*:*

$$\mathsf{P}(z) = [I_r, \mathbf{0}]\mathsf{P}_{J+1}(z)\mathsf{P}_J(z)\cdots\mathsf{P}_1(z)\mathsf{P}_0(z); \tag{3.7}$$

(3) $\mathsf{P}_j, 1 \leqslant j \leqslant J$ *are elementary:* $\mathsf{P}_j(z)\mathsf{P}_j^*(z) = I_s$ *and* $\operatorname{coeffsupp}(\mathsf{P}_j) \subseteq [-1, 1]$*;*

(4) $(\mathsf{P}_{j+1}, \mathsf{P}_j)$ *has mutually compatible symmetry for all* $0 \leqslant j \leqslant J$*;*

(5) $\mathsf{P}_0 = \mathsf{U}^*_{\mathcal{S}\theta_2}$ *and* $\mathsf{P}_{J+1} = diag(\mathsf{U}_{\mathcal{S}\theta_1}, I_{s-r})$*, where* $\mathsf{U}_{\mathcal{S}\theta_1}$*,* $\mathsf{U}_{\mathcal{S}\theta_2}$ *are products of a permutation matrix with a diagonal matrix of monomials, as defined in* (3.9)*;*

(6) $J \leqslant \max_{1\leqslant m\leqslant r, 1\leqslant n\leqslant s} \lceil|\mathrm{coeffsupp}([\mathsf{P}]_{m,n})|/2\rceil$*, where* $\lceil\cdot\rceil$ *is the ceiling function.*

The proof of Theorem 9 is given in [**?**], in which a general extension algorithm ($r \geqslant 1$) for the matrix extension with symmetry was also proposed. Since we are only concerning about the construction of tight d-wavelet frame in this paper, we shall provide an simple algorithm for the matrix extension with symmetry for the case $r = 1$.

Before we introduce our algorithm, we need some auxiliary results.

For a $1 \times n$ row vector $\mathtt{f}$ in $\mathbb{F}$ such that $\|\mathtt{f}\| \neq 0$, where $\|\mathtt{f}\|^2 := \mathtt{f}\mathtt{f}^*$, we define $n_{\mathtt{f}}$ to be the number of nonzero entries in $\mathtt{f}$ and $\mathbf{e}_j := [0, \dots, 0, 1, 0, \dots, 0]$ to be the jth unit coordinate row vector in $\mathbb{R}^n$. Let $E_{\mathtt{f}}$ be a permutation matrix such that $\mathtt{f}E_{\mathtt{f}} = [f_1, \dots, f_{n_{\mathtt{f}}}, 0, \dots, 0]$ with $f_j \neq 0$ for $j = 1, \dots, n_{\mathtt{f}}$. We define

$$V_{\mathtt{f}} := \begin{cases} \frac{\bar{f_1}}{|f_1|}, & \text{if } n_{\mathtt{f}} = 1; \\ \frac{f_1}{|f_1|}\left(I_n - \frac{2}{\|v_{\mathtt{f}}\|^2} v_{\mathtt{f}}^* v_{\mathtt{f}}\right), & \text{if } n_{\mathtt{f}} > 1, \end{cases} \tag{3.8}$$

where $v_{\mathtt{f}} := \mathtt{f} - \frac{f_1}{|f_1|}\|\mathtt{f}\|\mathbf{e}_1$. Observing that $\|v_{\mathtt{f}}\|^2 = 2\|\mathtt{f}\|(\|\mathtt{f}\| - |f_1|)$, we can verify that $V_{\mathtt{f}}V_{\mathtt{f}}^* = I_n$ and $\mathtt{f}E_{\mathtt{f}}V_{\mathtt{f}} = \|\mathtt{f}\|\mathbf{e}_1$. Let $U_{\mathtt{f}} := E_{\mathtt{f}}V_{\mathtt{f}}$. Then $U_{\mathtt{f}}$ is unitary and satisfies $U_{\mathtt{f}} = [\frac{\mathtt{f}^*}{\|\mathtt{f}\|}, F^*]$ for some $(n-1) \times n$ matrix F in $\mathbb{F}$ such that $\mathtt{f}U_{\mathtt{f}} = [\|\mathtt{f}\|, 0, \dots, 0]$. We also define $U_{\mathtt{f}} := I_n$ if $\mathtt{f} = \mathbf{0}$ and $U_{\mathtt{f}} := \emptyset$ if $\mathtt{f} = \emptyset$. Here, $U_{\mathtt{f}}$ plays the role of reducing the number of nonzero entries in $\mathtt{f}$.

Let θ be a $1 \times n$ row vector of Laurent polynomials with symmetry such that $\mathcal{S}\theta = [\varepsilon_1 z^{c_1}, \dots, \varepsilon_n z^{c_n}]$ for some $\varepsilon_1, \dots, \varepsilon_n \in \{-1, 1\}$ and $c_1, \dots, c_n \in \mathbb{Z}$. Then, the symmetry of any entry in the vector $\theta\mathrm{diag}(z^{-\lceil c_1/2\rceil}, \dots, z^{-\lceil c_n/2\rceil})$ belongs to $\{\pm 1, \pm z^{-1}\}$. Thus, there is a permutation matrix E_θ to regroup these four types of symmetries together so that

$$\mathcal{S}(\theta\mathsf{U}_{\mathcal{S}\theta}) = [\mathbf{1}_{n_1}, -\mathbf{1}_{n_2}, z^{-1}\mathbf{1}_{n_3}, -z^{-1}\mathbf{1}_{n_4}], \tag{3.9}$$

where $\mathsf{U}_{\mathcal{S}\theta} := \mathrm{diag}(z^{-\lceil c_1/2\rceil}, \dots, z^{-\lceil c_n/2\rceil})E_\theta$, $\mathbf{1}_m$ denotes the $1 \times m$ row vector $[1, \dots, 1]$, and $n_1, \dots, n_4$ are nonnegative integers uniquely determined by $\mathcal{S}\theta$. Note that $\mathsf{U}_{\mathcal{S}\theta}$ do not increase the length of the coefficient support of θ.

Let q be a $1 \times s$ row vector of Laurent polynomial satisfying $\mathsf{q}\mathsf{q}^* = 1$ and $\mathcal{S}\mathsf{q} = [\mathbf{1}_{s_1}, -\mathbf{1}_{s_2}, z^{-1}\mathbf{1}_{s_3}, -z^{-1}\mathbf{1}_{s_4}]$ for some nonnegative integers $s_1, \dots, s_4$ such that $s_1 + s_2 + s_3 + s_4 = s$. Then, q must take the form in (3.10) or (3.11) with $\mathtt{f}_1 \neq \mathbf{0}$ as follows:

$$\begin{aligned} \mathsf{q} = {} & [\mathtt{f}_1, -\mathtt{f}_2, \mathsf{g}_1, -\mathsf{g}_2]\mathsf{z}^{\ell_1} + [\mathtt{f}_3, -\mathtt{f}_4, \mathsf{g}_3, -\mathsf{g}_4]\mathsf{z}^{\ell_1+1} + \sum_{\ell=\ell_1+2}^{\ell_2-2} \mathrm{coeff}(\mathsf{q}, \ell)\mathsf{z}^{\ell} \\ & + [\mathtt{f}_3, \mathtt{f}_4, \mathsf{g}_1, \mathsf{g}_2]\mathsf{z}^{\ell_2-1} + [\mathtt{f}_1, \mathtt{f}_2, \mathbf{0}, \mathbf{0}]\mathsf{z}^{\ell_2}; \end{aligned} \tag{3.10}$$

$$\begin{aligned} \mathsf{q} = {} & [\mathbf{0}, \mathbf{0}, \mathtt{f}_1, -\mathtt{f}_2]\mathsf{z}^{\ell_1} + [\mathsf{g}_1, -\mathsf{g}_2, \mathtt{f}_3, -\mathtt{f}_4]\mathsf{z}^{\ell_1+1} + \sum_{\ell=\ell_1+2}^{\ell_2-2} \mathrm{coeff}(\mathsf{q}, \ell)\mathsf{z}^{\ell} \\ & + [\mathsf{g}_3, \mathsf{g}_4, \mathtt{f}_3, \mathtt{f}_4]\mathsf{z}^{\ell_2-1} + [\mathsf{g}_1, \mathsf{g}_2, \mathtt{f}_1, \mathtt{f}_2]\mathsf{z}^{\ell_2}. \end{aligned} \tag{3.11}$$

If q takes the form in (3.11), we further construct a permutation matrix E_{q} so that $[\mathsf{g}_1, \mathsf{g}_2, \mathtt{f}_1, \mathtt{f}_2]E_{\mathsf{q}} = [\mathtt{f}_1, \mathtt{f}_2, \mathsf{g}_1, \mathsf{g}_2]$ and define $\mathsf{U}_{\mathsf{q},\varepsilon} := E_{\mathsf{q}}\mathrm{diag}(I_{s-s_{\mathsf{g}}}, z^{-1}I_{s_{\mathsf{g}}})$, where s_{g} is the size of the row vector $[\mathsf{g}_1, \mathsf{g}_2]$. Then $\mathsf{q}\mathsf{U}_\mathsf{q}$ takes the form in (3.10). For $\mathsf{q}E_\varepsilon$ of form (3.10), we simply let $\mathsf{U}_\mathsf{q} := I_s$. In this way, $\mathsf{q}_0 := \mathsf{q}\mathsf{U}_\mathsf{q}$ always takes the form in (3.10) with $\mathtt{f}_1 \neq \mathbf{0}$.

Note that $\mathsf{U}_\mathsf{q}\mathsf{U}_\mathsf{q}^* = I_s$ and $\|\mathtt{f}_1\| = \|\mathtt{f}_2\|$ if $\mathsf{q}_0\mathsf{q}_0^* = 1$. Now an $s \times s$ paraunitary matrix $\mathsf{B}_{\mathsf{q}_0}$ to reduce the coefficient support of q_0 as in (3.10) with $\mathtt{f}_1 \neq \mathbf{0}$ from $[\ell_1, \ell_2]$ to $[\ell_1 + 1, \ell_2 - 1]$ is given by:

$$\mathsf{B}_{\mathsf{q}_0}^* := \frac{1}{c}\left[\begin{array}{c|c|c|c}
\mathtt{f}_1(z + \frac{c_0}{c_{\mathtt{f}_1}} + \frac{1}{z}) & \mathtt{f}_2(z - \frac{1}{z}) & \mathsf{g}_1(1 + \frac{1}{z}) & \mathsf{g}_2(1 - \frac{1}{z}) \\
cF_1 & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \hline
-\mathtt{f}_1(z - \frac{1}{z}) & -\mathtt{f}_2(z - \frac{c_0}{c_{\mathtt{f}_1}} + \frac{1}{z}) & -\mathsf{g}_1(1 - \frac{1}{z}) & -\mathsf{g}_2(1 + \frac{1}{z}) \\
\mathbf{0} & cF_2 & \mathbf{0} & \mathbf{0} \\ \hline
\frac{c_{\mathsf{g}_1}}{c_{\mathtt{f}_1}}\mathtt{f}_1(1 + z) & -\frac{c_{\mathsf{g}_1}}{c_{\mathtt{f}_1}}\mathtt{f}_2(1 - z) & c_{\mathsf{g}_1'}\mathsf{g}_1' & \mathbf{0} \\
\mathbf{0} & \mathbf{0} & cG_1 & \mathbf{0} \\ \hline
\frac{c_{\mathsf{g}_2}}{c_{\mathtt{f}_1}}\mathtt{f}_1(1 - z) & -\frac{c_{\mathsf{g}_2}}{c_{\mathtt{f}_1}}\mathtt{f}_2(1 + z) & \mathbf{0} & c_{\mathsf{g}_2'}\mathsf{g}_2' \\
\mathbf{0} & \mathbf{0} & \mathbf{0} & cG_2
\end{array}\right], \qquad (3.12)$$

where $c_{\mathtt{f}_1} := \|\mathtt{f}_1\|$, $c_{\mathsf{g}_1} := \|\mathsf{g}_1\|$, $c_{\mathsf{g}_2} := \|\mathsf{g}_2\|$, $c_0 := \mathrm{coeff}(\mathsf{q}_0, \ell_1 + 1)\mathrm{coeff}(\mathsf{q}_0^*, -\ell_2)/c_{\mathtt{f}_1}$,

$$\begin{aligned}
&c_{\mathsf{g}_1'} := \begin{cases} \frac{-2c_{\mathtt{f}_1} - \overline{c_0}}{c_{\mathsf{g}_1}} & \text{if } \mathsf{g}_1 \neq \mathbf{0}; \\ c & \text{otherwise,} \end{cases} \qquad c_{\mathsf{g}_2'} := \begin{cases} \frac{2c_{\mathtt{f}_1} - \overline{c_0}}{c_{\mathsf{g}_2}} & \text{if } \mathsf{g}_2 \neq \mathbf{0}; \\ c & \text{otherwise,} \end{cases} \\
&c := (4c_{\mathtt{f}_1}^2 + 2c_{\mathsf{g}_1}^2 + 2c_{\mathsf{g}_2}^2 + |c_0|^2)^{1/2},
\end{aligned} \qquad (3.13)$$

and $[\frac{\mathtt{f}_j^*}{\|\mathtt{f}_j\|}, F_j^*] = U_{\mathtt{f}_j}$, $[\mathsf{g}_j'^*, G_j^*] = U_{\mathsf{g}_j}$ are unitary constant extension matrices in $\mathbb{F}$ for vectors $\mathtt{f}_j, \mathsf{g}_j$ in $\mathbb{F}$, for $j = 1, 2$, respectively. The operations for the emptyset $\emptyset$ are defined by $\|\emptyset\| = \emptyset$, $\emptyset + A = A$ and $\emptyset \cdot A = \emptyset$ for any object A.

Lemma 10. *Let* q *be a* $1 \times s$ *row vector of Laurent polynomial satisfying* $\mathsf{q}\mathsf{q}^* = 1$, $|\mathrm{coeffsupp}(\mathsf{q})| > 2$, *and* $\mathcal{S}\mathsf{q} = [\mathbf{1}_{s_1}, -\mathbf{1}_{s_2}, z^{-1}\mathbf{1}_{s_3}, -z^{-1}\mathbf{1}_{s_4}]$ *for some nonnegative integers* $s_1, \ldots, s_4$ *such that* $s_1 + s_2 + s_3 + s_4 = s$. *Let* $\mathsf{B}_\mathsf{q} := \mathsf{U}_\mathsf{q}\mathsf{B}_{\mathsf{q}_0}\mathsf{U}_\mathsf{q}^*$ *with* U_q *and* $\mathsf{B}_{\mathsf{q}_0}$ *being constructed as above. Then,* $\mathcal{S}\mathsf{B}_\mathsf{q} = [\mathbf{1}_{s_1}, -\mathbf{1}_{s_2}, z\mathbf{1}_{s_3}, -z\mathbf{1}_{s_4}]^T[\mathbf{1}_{s_1}, -\mathbf{1}_{s_2}, z^{-1}\mathbf{1}_{s_3}, -z^{-1}\mathbf{1}_{s_4}]$, $\mathrm{coeffsupp}(\mathsf{B}_\mathsf{q}) = [-1, 1]$, *and* $\mathrm{coeffsupp}(\mathsf{q}\mathsf{B}_\mathsf{q}) = [\ell_1 + 1, \ell_2 - 1]$. *That is,* B_q *has compatible symmetry with coefficient support on* $[-1, 1]$ *and* B_q *reduces the length of the coefficient support of* q *exactly by* 2. *Moreover,* $\mathcal{S}(\mathsf{q}\mathsf{B}_\mathsf{q}) = \mathcal{S}\mathsf{q}$.

Proof. Direct computation shows that $\mathrm{coeff}(\mathsf{q}\mathsf{B}, k) = 0$ for $k \in \{\ell_1 - 1, \ell_1, \ell_2 + 1, \ell_2\}$. One can also show that $\mathrm{coeff}(\mathsf{q}\mathsf{B}, k) \neq 0$ for $k \in \{\ell_1 + 1, \ell_2 - 1\}$. Hence $\mathrm{coeffsupp}(\mathsf{q}\mathsf{B}_\mathsf{q}) = [\ell_1 + 1, \ell_2 - 1]$. The other parts of the Lemma follows from our construction. ■

Now, it is easy to prove Theorem 9 for $r = 1$ using Lemma 10. Moreover, we can have a constructive algorithm to derive P_e from a given column p, which mainly has three steps: initialization, support reduction, and finalization. The step of initialization reduces the symmetry pattern of p to a standard form. The step of support reduction is the main body of the algorithm, producing a sequence of elementary matrices

$\mathsf{A}_1, \dots, \mathsf{A}_J$ that reduce the length of the coefficient support of p to 0. The step of finalization generates the desired matrix P_e as in Theorem 9. More precisely, our algorithm written in the form of *pseudo-code* is as follows:

Algorithm 1. *Input* p*, which is a* $1 \times s$ *vector of Laurent polynomial with symmetry satisfying* $\mathsf{p}\mathsf{p}^* = 1$.
1. Initialization: *Let* $\mathsf{q} := \mathsf{p}\mathsf{U}_{\mathcal{S}\mathsf{p}}$. *Then* $\mathcal{S}\mathsf{q} = [\mathbf{1}_{s_1}, -\mathbf{1}_{s_2}, z^{-1}\mathbf{1}_{s_3}, -z^{-1}\mathbf{1}_{s_4}]$*, where all nonnegative integers* $r_1, \dots, r_4, s_1, \dots, s_4$ *are uniquely determined by* $\mathcal{S}\mathsf{p}$.
2. Support Reduction: *Let* $\mathsf{P}_0 := \mathsf{U}^*_{\mathcal{S}\mathsf{p}}$ *and* $J := 1$.

```
while (|coeffsupp($\mathsf{q}$)| > 0) do
  if |coeffsupp($\mathsf{q}$)| > 1 then
    $\mathsf{q} := \mathsf{q}\mathsf{B}_\mathsf{q}$, $\mathsf{A}_J := \mathsf{B}_\mathsf{q}$, and $\mathsf{P}_J := \mathsf{A}^*_J$.
  else
    $\mathsf{q}$ is of the form: $\mathsf{q} = [\mathbf{0}, \mathbf{0}, \mathsf{g}_1, -\mathsf{g}_2]z^{-1} + [\mathtt{f}_3, \mathbf{0}, \mathsf{g}_1, \mathsf{g}_2]$; Let
        $\mathsf{B}_J := \left[\begin{array}{c|cc|cc} I_{s_1+s_2} & & & & \\ \hline & \frac{\mathsf{g}_1^*(1+z)}{2\|\mathsf{g}_1\|} & G_1 & \frac{\mathsf{g}_1^*(1-z)}{2\|\mathsf{g}_1\|} & \mathbf{0} \\ \hline & \frac{\mathsf{g}_2^*(1-z)}{2\|\mathsf{g}_1\|} & \mathbf{0} & \frac{\mathsf{g}_2^*(1+z)}{2\|\mathsf{g}_1\|} & G_2 \end{array}\right]$,
    where $[\frac{\mathsf{g}_j^*}{\|\mathsf{g}_j\|}, G_j] = U_{\mathsf{g}_j}$, $j = 1, 2$. $\mathsf{B}_J$ is paraunitary, |coeffsupp($\mathsf{q}\mathsf{B}_J$)| = 0, and
            $\mathcal{S}(\mathsf{q}\mathsf{B}_J) = [\mathbf{1}_{s_1}, -\mathbf{1}_{s_2}, 1, z^{-1}\mathbf{1}_{s_3-1}, -1, -z^{-1}\mathbf{1}_{s_4-1}]$.
    Let $E$ be a permutation matrix such that
            $\mathcal{S}(\mathsf{q}\mathsf{B}_J)E = [\mathbf{1}_{s_1+1}, -\mathbf{1}_{s_2+1}, z^{-1}\mathbf{1}_{s_3-1}, -z^{-1}\mathbf{1}_{s_4-1}]$.
    Let $\mathsf{A}_J := \mathsf{B}_J E$, $\mathsf{P}_J := \mathsf{A}^*_J$, and $\mathsf{q} = \mathsf{q}\mathsf{A}_J$.
  end if
  $J := J + 1$
end while
```

3. Finalization: $\mathsf{q} = [\mathtt{f}, \mathbf{0}, \mathbf{0}, \mathbf{0}]$ *for some* $1 \times s_1'$ *constant vectors* $\mathtt{f}$ *in* $\mathbb{F}$. *Let* $U := \mathrm{diag}(U_\mathtt{f}, I_{s-s_1'})$. *Define* $\mathsf{P}_J := U^*_\mathtt{f}$.
Output a desired matrix P_e *satisfying all the properties in Theorem 9 for* $r = 1$.

4. Symmetric Complex Tight d-frame via Matrix Extension

In this section, we shall discuss the application of our results on matrix extension with symmetry to d-band symmetric paraunitary filter banks in electronic engineering and to orthonormal multiwavelets with symmetry in wavelet analysis. In order to do so, let us introduce some definitions first.

Let $\mathbb{F}$ be a subfield of $\mathbb{C}$ such that (3.6) holds. Let ϕ be a compactly supported d-refinable function in $L_2(\mathbb{R})$ associated with a low-pass filter $a_0 : \mathbb{Z} \mapsto \mathbb{F}$. Recall that its symbol is defined to be $\mathsf{a}_0(z) := \sum_{k\in\mathbb{Z}} a_0(k)z^k$, which is a Laurent polynomials with coefficients in $\mathbb{F}$. We denote its d-*band subsymbols* by $\mathsf{a}_{0;\gamma}(z) := \sqrt{\mathsf{d}} \sum_{k\in\mathbb{Z}} a_0(\gamma + \mathsf{d}k)z^k$, $\gamma = 0, \dots, \mathsf{d} - 1$. To construct a tight d-wavelet frame from ϕ, one has to design high-pass filters $a^1, \dots, a^L : \mathbb{Z} \mapsto \mathbb{F}^{r\times r}$ such that (1.7) holds. Let $\mathcal{P}(z)$ be the polyphase matrix defined by

$$\mathcal{P}(z) := \begin{bmatrix} \mathsf{a}_{0;0}(z) & \cdots & \mathsf{a}_{0;\mathsf{d}-1}(z) \\ \mathsf{a}_{1;0}(z) & \cdots & \mathsf{a}_{1;\mathsf{d}-1}(z) \\ \vdots & \vdots & \vdots \\ \mathsf{a}_{L;0}(z) & \cdots & \mathsf{a}_{L;\mathsf{d}-1}(z) \end{bmatrix}. \tag{4.1}$$

Then it is easily seen that (1.7) is equivalent to

$$\mathcal{P}(z)^* \mathcal{P}(z) = I_{\mathsf{d}}, \tag{4.2}$$

where each $\mathsf{a}_{m;\gamma}$ is a subsymbol of a^m for $m = 1, \ldots, L; \gamma = 0, \ldots, \mathsf{d} - 1$, respectively.

Symmetry of the filters is a very much desirable property in many applications. We say that the low-pass filter a_0 (or a_0) has symmetry if

$$\mathsf{a}_0(z) = z^{(\mathsf{d}-1)c_0} \mathsf{a}_0(1/z) \tag{4.3}$$

for some $c_0 \in \mathbb{R}$ such that $(\mathsf{d} - 1)c_0 \in \mathbb{Z}$. If a_0 has symmetry as in (4.3) and if 1 is a simple eigenvalue of $\mathsf{a}_0(1)$, then it is well known that the d-refinable function ϕ associated with the low-pass filter a_0 has symmetry: $\phi(c_0 - \cdot) = \phi$.

Under the symmetry condition in (4.3),

$$\mathsf{a}_{0;\gamma}(z) = z^{R_\gamma} \mathsf{a}_{0;Q_\gamma}(z^{-1}) \quad \gamma = 0, \ldots, \mathsf{d} - 1, \tag{4.4}$$

where $\gamma, Q_\gamma \in \Gamma := \{0, \ldots, \mathsf{d} - 1\}$ and R_γ, Q_γ are uniquely determined by

$$(\mathsf{d} - 1)c_0 - \gamma = \mathsf{d}R_\gamma + Q_\gamma \quad \text{with} \quad R_\gamma \in \mathbb{Z},\ Q_\gamma \in \Gamma. \tag{4.5}$$

Now, we can easily deduce symmetric Laurent polynomial via (4.4). In fact, let

$$\mathsf{b}_{0;\gamma}(z) := \begin{cases} \mathsf{a}_{0;\gamma}(z), & \gamma = Q_\gamma; \\ \frac{1}{\sqrt{2}}(\mathsf{a}_{0;\gamma}(z) + z^{\kappa_\gamma} \mathsf{a}_{0;Q_\gamma}(z)), & \gamma < Q_\gamma; \\ \frac{1}{\sqrt{2}}(\mathsf{a}_{0;\gamma}(z) - z^{\kappa_\gamma} \mathsf{a}_{0;Q_\gamma}(z)), & \gamma > Q_\gamma, \end{cases} \tag{4.6}$$

where $\gamma = 0, \ldots, \mathsf{d} - 1$ and each κ_γ is an integer so that the length of the coefficient support of $\mathsf{a}_{0;\gamma}(z) + z^{\kappa_\gamma} \mathsf{a}_{0;Q_\gamma}(z)$ is minimal. Then, one can show that each $\mathsf{b}_{0;\gamma}(z)$ is of a Laurent polynomial with symmetry. Note that the transform matrix U with respect to (4.6) is paraunitary.

Next, we show that we can construct a vector of Laurent polynomial with symmetry from a low-pass filter a for the complex pseudo spline of type I with order (m, n) to which Algorithm 1 is applicable.

Fixed $m, n \in \mathbb{N}$ such that $1 \leqslant 2n - 1 \leqslant m$. For simplicity, let $a_0 = a$ and a_0 be its symbol. From (4.6), we can construct a $1 \times \mathsf{d}$ vector of Laurent polynomial $\mathsf{p}(z) = [\mathsf{b}_{0;0}(z), \ldots, \mathsf{b}_{0;\mathsf{d}-1}(z)]$ from a_0. However, $\mathsf{p}\mathsf{p}^* \neq 1$ when $n < m$. To apply our matrix extension algorithm, we need to append extra entries to p. It is easy to show that

$$1 - \sum_{j=0}^{\mathsf{d}-1} |\widehat{a_0}(\xi + 2j\pi/\mathsf{d})|^2 = 1 - \sum_{\gamma=0}^{\mathsf{d}-1} \mathsf{a}_{0;\gamma}(z^{\mathsf{d}}) \mathsf{a}^*_{0;\gamma}(z^{\mathsf{d}}), \quad z = e^{-i\xi},$$

where $\mathsf{a}_{0;\gamma}, \gamma = 0, \ldots, \mathsf{d}-1$ are the subsymbols of a_0. By Corollary 5, we have

$$1 - \sum_{j=0}^{\mathsf{d}-1} |\widehat{a}_0(\xi + 2j\pi/\mathsf{d})|^2 = |\widehat{b}_0(\mathsf{d}\xi)|^2.$$

for some 2π-periodic trigonometric function $\widehat{b}_0$ with real coefficients. Hence, we can construct a Laurent polynomial $\mathsf{a}_{0;\mathsf{d}}(z)$ from $\widehat{b}_0$ such that $\mathsf{a}_{0;\mathsf{d}}(e^{-i\xi}) = \widehat{b}_0(\xi)$. Then, the vector of Laurent polynomials $\mathsf{q}(z) = [\mathsf{a}_{0;0}(z), \ldots, \mathsf{a}_{0;\mathsf{d}-1}(z), \mathsf{a}_{0;\mathsf{d}}(z)]$ satisfies $\mathsf{q}\mathsf{q}^* = 1$.

For $m = 2n$, by Corollary 5, $|\widehat{b}_0(\xi)|^2 = c_{2n,2n-1}[\sin^2(\xi/2)/\mathsf{d}^2]^{2n-1}$. W $\mathsf{a}_{0;\mathsf{d}}(z)$ can be constructed explicitly as follows:

$$\mathsf{a}_{0;\mathsf{d}}(z) = \sqrt{c_{2n,2n-1}} \left(\frac{2 - z - 1/z}{4\mathsf{d}^2}\right)^{n-1} \frac{1-z}{2\mathsf{d}}.$$

Then $\mathsf{a}_{0;\mathsf{d}}(z)$ is symmetric, let $\mathsf{b}_{0;\mathsf{d}}(z) = \mathsf{a}_{0;\mathsf{d}}(z)$. Then $\mathsf{p} := [\mathsf{b}_{0;0}, \ldots, \mathsf{b}_{0;\mathsf{d}}]$ is a $1 \times (\mathsf{d}+1)$ vector of Laurent polynomials with symmetry satisfying $\mathsf{p}\mathsf{p}^* = 1$.

If $m \neq 2n$, $\mathsf{a}_{0;\mathsf{d}}(z)$ is a not necessary symmetric, we can further let $\mathsf{b}_{0;\mathsf{d}}(z) = (\mathsf{a}_{0;\mathsf{d}}(z) + \mathsf{a}_{0;\mathsf{d}}(1/z))/2$ and $\mathsf{b}_{0;\mathsf{d}+1}(z) = (\mathsf{a}_{0;\mathsf{d}}(z) - \mathsf{a}_{0;\mathsf{d}}(1/z))/2$. In this way, $\mathsf{p} := [\mathsf{b}_{0;0}, \ldots, \mathsf{b}_{0;\mathsf{d}}, \mathsf{b}_{0;\mathsf{d}+1}]$ is a $1 \times (\mathsf{d}+2)$ vector of Laurent polynomials with symmetry satisfying $\mathsf{p}\mathsf{p}^* = 1$.

Consequently, we can summerize the above result as follows:

Theorem 11. *Let $m, n \in \mathbb{N}$ be such that $1 \leqslant n \leqslant m$. Let a_0 be the low-pass filter for the complex pseudo spline of order (m, n) defined in* (1.12) *and $\mathsf{a}_0(z)$ be its symbol. Then, one can derive Laurent polynomials $\mathsf{a}_{0;\mathsf{d}}(z), \ldots, \mathsf{a}_{0;L}(z), \mathsf{d} \leqslant L \leqslant \mathsf{d}+1$ such that $\mathsf{p}_{\mathsf{a}_0} := [\mathsf{a}_{0;0}, \ldots, \mathsf{a}_{0;\mathsf{d}-1}, \cdots, \mathsf{a}_{0;L}(z)]$ satisfies $\mathsf{p}_{\mathsf{a}_0}\mathsf{p}_{\mathsf{a}_0}^* = 1$, where $\mathsf{a}_{0;0}, \ldots, \mathsf{a}_{0;\mathsf{d}-1}$ are subsymbols of a_0. Moreover, one can construct a $(L+1) \times (L+1)$ paraunitary matrix U such that $\mathsf{p}_{\mathsf{a}_0}\mathsf{U}$ is a vector of Laurent polynomial with symmetry. In particular, for $m = 2n$, $\mathsf{a}_{0;\mathsf{d}}(z) = \sqrt{c_{2n,2n-1}}\left(\frac{2-z-1/z}{4\mathsf{d}^2}\right)^{n-1} \frac{1-z}{2}$ and $L = \mathsf{d}$.*

Now, applying Theorem 9, we have the following algorithm to construct high-pass filters $\mathsf{a}_1, \ldots, \mathsf{a}_L$ from a low-pass filter a_0 for a complex pseudo spline of type I with order (m, n).

Algorithm 2. *Input low-pass filter a_0 for a complex pseudo spline of type I with order (m, n), $1 \leqslant 2n - 1 \leqslant m$. a_0 satisfies* (4.3)*.*

(1) *Construct $\mathsf{p}_{\mathsf{a}_0}(z)$ and U as in Theorem 11 such that $\mathsf{p} := \mathsf{p}_{\mathsf{a}_0}\mathsf{U}$ is a $1 \times (L+1)$ row vector of Laurent polynomials with symmetry ($L = \mathsf{d}$ when $m = 2n$ while $L = \mathsf{d}+1$ when $m \neq 2n$.*

(2) *Derive P_e with all the properties as in Theorem 9 for the case $r = 1$ from p by Algorithm 1.*

(3) *Let $\mathcal{P} := \mathsf{P}_e\mathsf{U}^* =: (\mathsf{a}_{m;\gamma})_{0 \leqslant m \leqslant L, 0 \leqslant \gamma \leqslant \mathsf{d}-1}$ as in* (4.1)*. Define high-pass filters*

$$\mathsf{a}_m(z) := \frac{1}{\sqrt{\mathsf{d}}} \sum_{\gamma=0}^{\mathsf{d}-1} \mathsf{a}_{m;\gamma}(z^{\mathsf{d}}) z^{\gamma}, \qquad m = 1, \ldots, L. \tag{4.7}$$

Note that we only need the first d columns of $\mathsf{P}_e\mathsf{U}^*$.

Output a symmetric filter bank $\{\mathsf{a}_0, \mathsf{a}_1, \dots, \mathsf{a}_L\}$ *with the perfect reconstruction property, i.e.* $\mathcal{P}^*(z)\mathcal{P}(z) = I_{\mathsf{d}}$*. All filters* a_m*,* $m = 1, \dots, \mathsf{d}-1$*, have symmetry:*

$$\mathsf{a}_m(z) = \varepsilon_m z^{\mathsf{d}c_m - c_0} \mathsf{a}_m(1/z), \tag{4.8}$$

where $c^m := k_m + c_0 \in \mathbb{R}$ *and all* $\varepsilon_m \in \{-1, 1\}$*,* $k_m \in \mathbb{Z}$ *for* $m = 1, \dots, L$ *are determined by the symmetry pattern of* P_e *as follows:*

$$[1, \varepsilon_1 z^{k_1}, \dots, \varepsilon_L z^{k_L}]^T \mathcal{S}\mathsf{p} := \mathcal{S}\mathsf{P}_e. \tag{4.9}$$

Proof. Rewrite $\mathsf{P}_e = (\mathsf{b}_{m;\gamma})_{0 \leqslant m, \gamma \leqslant L}$. Since P_e has compatible symmetry as in (4.9), we have $\mathcal{S}\mathsf{b}_{m;\gamma} = \varepsilon_m z_m^k \mathcal{S}\mathsf{b}_{0;\gamma}$ for $m = 1, \dots, \mathsf{d}-1$. By (4.6), we have $\mathcal{S}\mathsf{b}_{0;\gamma} = \operatorname{sgn}(Q_\gamma - \gamma) z^{R_\gamma + k_\gamma}$, $\gamma = 0, \dots, \mathsf{d}-1$, where $\operatorname{sgn}(x) = 1$ for $x \geqslant 0$ and $\operatorname{sgn}(x) = -1$ for $x < 0$. Then, we have

$$\mathcal{S}\mathsf{b}_{m;\gamma} = \operatorname{sgn}(Q_\gamma - \gamma)\varepsilon_m z^{R_\gamma + \kappa_\gamma + k_m}, \tag{4.10}$$

By (4.10) and the transformation matrix U^* with respect to (4.6), we deduce that

$$\mathsf{a}_{m;\gamma} = \varepsilon_m z^{R_\gamma + k_m} \mathsf{a}_{m;Q_\gamma}(z^{-1}). \tag{4.11}$$

This implies that $\mathcal{S}\mathsf{a}_m = \varepsilon_m z^{\mathsf{d}(k_m + c_0) - c_0}$, which is equivalent to (4.8) with $c_m := k_m + c_0$ for $m = 1, \dots, L$.

Since the high-pass filters $\mathsf{a}_1, \dots, \mathsf{a}_L$ satisfy (4.8), it is easy to verify that $\psi^1, \dots, \psi^L$ defined in (1.4) also has the following symmetry:

$$\psi^1(c_1 - \cdot) = \varepsilon_1 \psi^1, \quad \psi^2(c_2 - \cdot) = \varepsilon_2 \psi^2, \quad \dots, \quad \psi^L(c_L - \cdot) = \varepsilon_L \psi^1, . \tag{4.12}$$

In fact, by (4.8), (1.4) and the symmetry of ϕ, we have

$$\begin{aligned}\widehat{\psi^m}(-\mathsf{d}\xi) &= \mathsf{a}_m(e^{i\xi})\widehat{\phi}(-\xi) = \varepsilon_m e^{-i(\mathsf{d}c_m - c_0)} \mathsf{a}_m(e^{-i\xi}) e^{-ic_0\xi} \widehat{\phi}(\xi) \\ &= \varepsilon_m e^{-i(\mathsf{d}c_m)\xi} \mathsf{a}_m(e^{-i\xi})\widehat{\phi}(\xi) = \varepsilon_m e^{-i(\mathsf{d}c_m)\xi} \widehat{\psi^m}(\mathsf{d}\xi).\end{aligned}$$

This implies $\widehat{\psi^m}(-\xi) = \varepsilon_m e^{-ic_m\xi} \widehat{\psi^m}(\xi)$, which is equivalent to (4.12).

In the following, let us present some examples to demonstrate our results and illustrate our algorithms.

Example 1. Consider dilation factor $\mathsf{d} = 2$. Let $m = 4, n = 2$. Then $P_{4,3}(y) = 1 + 4y + 4y^2$. The low-pass filter $a_0^{4,2}$ with its symbol a_0 for the complex pseudo spline of order $(4, 2)$ is given by

$$\mathsf{a}_0(z) = \left(\frac{1+z}{2}\right)^4 \left[\left(-\frac{1}{2} - \frac{\sqrt{6}}{4}i\right)\frac{1}{z} + \left(2 + \frac{\sqrt{6}}{2}i\right)\frac{1}{z^2} + \left(-\frac{1}{2} - \frac{\sqrt{6}}{4}i\right)\frac{1}{z^3}\right].$$

Note that $\operatorname{coeffsupp}(\mathsf{a}_0) = [-3, 3]$ and $\mathsf{a}(z) = \mathsf{a}(z^{-1})$. In this case, $m = 2n$, by Theorem 11, we can obtain $\mathsf{p}_{\mathsf{a}_0} = [\mathsf{a}_{0;0}(z), \mathsf{a}_{0;1}(z), \mathsf{a}_{0;2}(z)]$ as follows:

$$\begin{aligned}\mathsf{a}_{0;0}(z) &= -\frac{\sqrt{3}i}{48}\left(3z + 8\sqrt{6}i - 6 + 3z^{-1}\right); \\ \mathsf{a}_{0;1}(z) &= -\frac{(\sqrt{2} + \sqrt{3}i)}{160}(5z + 8\sqrt{6}i - 26 + 5z^{-1})(1 + z^{-1}); \\ \mathsf{a}_{0;2}(z) &= -\frac{\sqrt{5}}{32}(z - 2 + z^{-1})(1 - z^{-1}).\end{aligned}$$

$\mathsf{p}_{\mathsf{a}_0}$ is already a 1×3 vector of Laurent polynomials with symmetry, i.e., $\mathcal{S}\mathsf{p}_{\mathsf{a}_0} = [1, z^{-1}, -z^{-1}]$. Applying Algorithm 2, we can obtain its 3×3 extension matrix $\mathsf{P}_e^* = [\mathsf{p}^*_{\mathsf{a}_0}, \mathsf{p}^*_{\mathsf{a}_1}, \mathsf{p}^*_{\mathsf{a}_2}]$ with $\mathsf{p}_{\mathsf{a}_1} := [\mathsf{a}_{1;0}, \mathsf{a}_{1;1}, \mathsf{a}_{1;2}]$ and $\mathsf{p}_{\mathsf{a}_2} := [\mathsf{a}_{2;0}, \mathsf{a}_{2;1}, \mathsf{a}_{2;2}]$ as follows:

$$\begin{aligned}
\mathsf{a}_{1;0}(z) &= \frac{-\sqrt{35}(\sqrt{6}+3i)}{1680}\left(3z - 8\sqrt{6}i + 42 + 3z^{-1}\right);\\
\mathsf{a}_{1;1}(z) &= \frac{-\sqrt{35}(2\sqrt{6}+i)}{5600}\left(5z - 8\sqrt{6}i - 74 + 5z^{-1}\right)(1+z^{-1});\\
\mathsf{a}_{1;2}(z) &= \frac{-\sqrt{7}(\sqrt{3}-\sqrt{2}i)}{1120}\left(5z - 16\sqrt{6}i - 58 + 5z^{-1}\right)(1-z^{-1});\\
\mathsf{a}_{2;0}(z) &= \frac{\sqrt{42}i}{56}(z - z^{-1});\\
\mathsf{a}_{2;1}(z) &= \frac{\sqrt{7}(2+\sqrt{6}i)}{560}\left(5z - 4\sqrt{6}i - 22 + 5z^{-1}\right)(1-z^{-1});\\
\mathsf{a}_{2;2}(z) &= \frac{\sqrt{70}(2\sqrt{6}+i)}{560}\left(5z - 12\sqrt{6}i - 26 + 5z^{-1}\right)(1+z^{-1}).
\end{aligned}$$

Note that the symmetry of P_e satisfies $\mathcal{S}\mathsf{P}_e = [1, 1, -1]^T \mathcal{S}\mathsf{p}_{\mathsf{a}_0}$ and the coefficient support of P_e satisfies $\mathrm{coeffsupp}([\mathsf{P}_e]_{:,j}) \subseteq \mathrm{coeffsupp}([\mathsf{p}_{\mathsf{a}_0}]_j)$ for $j = 1, 2, 3$. The high-pass filters $\mathsf{a}_1, \mathsf{a}_2$ constructed from $\mathsf{p}_{\mathsf{a}_1}$ and $\mathsf{p}_{\mathsf{a}_2}$ via (4.7) are then given by:

$$\begin{aligned}
\mathsf{a}_1(z) &= -\frac{\sqrt{35}(4\sqrt{3}+\sqrt{2}i)}{11200}(5z + 26 + 2\sqrt{6}i + 5z^{-1})(z - 2 + z^{-1})^2;\\
\mathsf{a}_2(z) &= \frac{\sqrt{7}(\sqrt{2}+\sqrt{3}i)}{560}(5z + 16 + 2\sqrt{6}i + 5z^{-1})(z - 2 + z^{-1})(z - z^{-1}).
\end{aligned}$$

We have $\mathsf{a}_1(z) = \mathsf{a}_1(z^{-1})$ and $\mathsf{a}_2(z) = -\mathsf{a}_2(z^{-1})$. Let ϕ be the 2-refinable function associated with the low-pass filter a_0. Let ψ^1, ψ^2 be the wavelet functions associated with the high-pass filters $\mathsf{a}_1, \mathsf{a}_2$ respectively by (1.4). Then $\phi(-\cdot) = \phi$, $\psi^1(-\cdot) = \psi^1$ and $\psi^2(-\cdot) = -\psi^2$. The graphs of ϕ, ψ^1, ψ^2 are as follows:



Figure 4.1: The graphs of ϕ, ψ^1, ψ^2 (left to right). Real part:solid line. Imaginary part:dashed line.

Example 2. Consider dilation factor $\mathsf{d} = 3$. Let $m = 4, n = 2$. Then $P_{4,3}(y) = 1 + \frac{32}{3}y + 64y^2)$. The low-pass filter $a_0^{4,2}$ with its symbol a_0 for the complex pseudo spline of order $(4, 2)$ is given by

$$\mathsf{a}_0(z) = \left(\frac{\frac{1}{z} + 1 + z}{3}\right)^4 \left[\left(-\frac{4}{3} - \frac{2\sqrt{5}}{3}i\right)\frac{1}{z} + \left(\frac{11}{3} + \frac{4\sqrt{5}}{3}i\right) + \left(-\frac{4}{3} - \frac{2\sqrt{5}}{3}i\right)z\right].$$

Note that $\operatorname{coeffsupp}(\mathsf{a}_0) = [-5, 5]$ and $\mathsf{a}(z) = \mathsf{a}(z^{-1})$. In this case, $m = 2n$, by Theorem 11, we can obtain $\mathsf{p}_{\mathsf{a}_0} = [\mathsf{a}_{0;0}(z), \mathsf{a}_{0;1}(z), \mathsf{a}_{0;2}(z), \mathsf{a}_{0;3}(z)]$ as follows:

$$\begin{aligned}
\mathsf{a}_{0;0}(z) &= -\frac{\sqrt{15}i}{405}\left(10z + 27\sqrt{5}i - 20 + 10z^{-1}\right);\\
\mathsf{a}_{0;1}(z) &= \frac{\sqrt{3}}{243}(-(4 + 2\sqrt{5}i)z^{-2} + 30z^{-1} + 60 + 6\sqrt{5}i - (5 + 4\sqrt{5}i)z);\\
\mathsf{a}_{0;2}(z) &= \frac{\sqrt{3}}{243}(-(5 + 4\sqrt{5}i)z^{-2} + (60 + 6\sqrt{5}i)z^{-1} + 30 - (4 + 2\sqrt{5}i)z)\\
\mathsf{a}_{0;3}(z) &= -\frac{2\sqrt{10}}{81}(z - 2 + z^{-1})(1 - z).
\end{aligned}$$

Note that $\mathsf{a}_{0;1}(z) = z^{-1}\mathsf{a}_{0;2}(z^{-1})$. Let $\mathsf{p} = \mathsf{p}_{\mathsf{a}_0}\mathsf{U}$ with U being the paraunitary matrix given by

$$\mathsf{U} := \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 0 & z^{-1} \end{bmatrix}.$$

Then p is a 1×4 vector of Laurent polynomials with symmetry pattern satisfying $\mathcal{S}\mathsf{p} = [1, z^{-1}, -z^{-1}, -z^{-1}]$. Applying Algorithm 2, we can obtain a 4×4 extension matrix $\mathsf{P}_e^* = [\mathsf{p}_{\mathsf{a}_0}^*, \mathsf{p}_{\mathsf{a}_1}^*, \mathsf{p}_{\mathsf{a}_2}^*, \mathsf{p}_{\mathsf{a}_3}]$ with $\mathsf{p}_{\mathsf{a}_1} := [\mathsf{a}_{1;0}, \mathsf{a}_{1;1}, \mathsf{a}_{1;2}, \mathsf{a}_{1;3}]$, $\mathsf{p}_{\mathsf{a}_2} := [\mathsf{a}_{2;0}, \mathsf{a}_{2;1}, \mathsf{a}_{2;2}, \mathsf{a}_{2;3}]$, and $\mathsf{p}_{\mathsf{a}_3} := [\mathsf{a}_{3;0}, \mathsf{a}_{3;1}, \mathsf{a}_{3;2}, \mathsf{a}_{3;3}]$. The coefficient support of P_e satisfies $\operatorname{coeffsupp}([\mathsf{P}_e]_{:,j}) \subseteq \operatorname{coeffsupp}([\mathsf{p}_{\mathsf{a}_0}]_j)$ for $j = 1, 2, 3, 4$. The high-pass filters $\mathsf{a}_1, \mathsf{a}_2, \mathsf{a}_3$ constructed from $\mathsf{p}_{\mathsf{a}_1}$, $\mathsf{p}_{\mathsf{a}_2}$, and $\mathsf{p}_{\mathsf{a}_3}$ via (4.7) are then given by:

$$\begin{aligned}
\mathsf{a}_1(z) &= \frac{\sqrt{19178}}{4660254}(b_1(z) + b_1(z^{-1}));\\
\mathsf{a}_2(z) &= \frac{\sqrt{218094}}{17665614}(b_2(z) - b_2(z^{-1}));\\
\mathsf{a}_3(z) &= \frac{2\sqrt{1338}}{54189}(b_3(z) - z^3 b_3(z^{-1})).
\end{aligned}$$

where

$$\begin{aligned}
b_1(z) = {} & \left(-172 - 86\,i\sqrt{5}\right)z^5 + \left(-215 - 172\,i\sqrt{5}\right)z^4 - 258\,i\sqrt{5}z^3\\
& + \left(1470 + 1224\,i\sqrt{5}\right)z^2 + \left(1860 + 2328\,i\sqrt{5}\right)z - 3036\,i\sqrt{5} - 2943\\
b_2(z) = {} & \left(-652 - 326\,i\sqrt{5}\right)z^5 + \left(-815 - 652\,i\sqrt{5}\right)z^4 - 978\,i\sqrt{5}z^3\\
& + \left(1832\,i\sqrt{5} + 1750\right)z^2 + \left(3508\,i\sqrt{5} + 3020\right)z\\
b_3(z) = {} & \left(4\sqrt{5} + 10\,i\right)z^5 + \left(5\sqrt{5} + 20\,i\right)z^4 + 30\,iz^3 + \left(-53\sqrt{5} - 260\,i\right)z^2.
\end{aligned}$$

We have $\mathsf{a}_1(z) = \mathsf{a}_1(z^{-1})$, $\mathsf{a}_2(z) = -\mathsf{a}_2(z^{-1})$, and $\mathsf{a}_3(z) = -z^3\mathsf{a}_3(z^{-1})$. Let ϕ be the 3-refinable function associated with the low-pass filter a_0. Let ψ^1, ψ^2, ψ^3 be the wavelet functions associated with the high-pass filters $\mathsf{a}_1, \mathsf{a}_2, \mathsf{a}_3$ respectively by (1.4). Then



Figure 4.2: The graphs of $\phi, \psi^1, \psi^2, \psi^3$ (left to right). Real part:solid line. Imaginary part:dashed line.

$\phi(-\cdot) = \phi$, $\psi^1(-\cdot) = \psi^1$, $\psi^2(-\cdot) = -\psi^2$, and $\psi^3(1-\cdot) = -\psi^3$. See Figure 2 for the graphs of ϕ, ψ^1, ψ^2, and ψ^3.

Example 3. Consider dilation factor $\mathsf{d} = 3$. Let $m = 5, n = 2$. Then $P_{5,3}(y) = 1 + \frac{40}{3}y + \frac{880}{9}y^2)$. The low-pass filter $a_0^{5,2}$ with its symbol $\mathsf{a}_0^{5,2}$ for the complex pseudo spline of order $(5,2)$ is given by

$$\mathsf{a}_0(z) = \left(\frac{\frac{1}{z} + 1 + z}{3}\right)^5 \left[\left(-\frac{5}{3} - \frac{\sqrt{30}}{3}i\right)\frac{1}{z} + \left(\frac{13}{3} + \frac{2\sqrt{30}}{3}i\right) + \left(-\frac{5}{3} - \frac{\sqrt{30}}{3}i\right)z\right].$$

In this case, $m = 2n - 1$, we have

$$\begin{aligned} 1 - \sum_{\gamma=0}^{2} \mathsf{a}_{0;\gamma}(z)^* \mathsf{a}_{0;\gamma}(z) = \frac{1}{19683}&(1-z)(1-z^{-1})(4z^3 + 106z^2 - 541z \\ &+ 2320 - 541z^{-1} + 106z^{-2} + 4z^{-3}), \quad z = e^{-i\xi}. \end{aligned}$$

By Riesz lemma, we can factorize the above positive trigonometric function and obtain $\mathsf{a}_{0;\mathsf{d}}(z) = c_0(z - r_1)(z^2 - r_2 z + r_3)(1 - z)$ with $r_1 = c_1 + \sqrt{c_1^2 - 1}$, $r_2 = c_2 + \overline{c_2}$ and $r_3 = |c_2|^2$, where $c_1 \in \mathbb{R}, c_2, \overline{c_2} \in \mathbb{C}$ are the roots for the polynomial $16y^3 + 212y^2 - 553y + 1054$ and $c_0 = \frac{2\sqrt{3}}{243\sqrt{-r_1 r_3}} \in \mathbb{R}$.

Let $\mathsf{p}_{\mathsf{a}_0} = [\mathsf{a}_{0;0}(z), \mathsf{a}_{0;1}(z), \mathsf{a}_{0;2}(z), z^{-2}\frac{1}{\sqrt{2}}\mathsf{a}_{0;3}(z), z^{-2}\frac{1}{\sqrt{2}}\mathsf{a}_{0;3}(z^{-1})]$. Then $\mathsf{p}_{\mathsf{a}_0}\mathsf{p}_{\mathsf{a}_0}^* = 1$. Note that $\mathsf{a}_{0;1}(z) = z^{-1}\mathsf{a}_{0;2}(z^{-1})$. Let $\mathsf{p} = \mathsf{p}_{\mathsf{a}_0}\mathsf{U}$ with U being the paraunitary matrix given by

$$\mathsf{U} := \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & 0 & 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix}.$$

Then p is a 1×5 vector of Laurent polynomials with symmetry pattern satisfying $\mathcal{S}\mathsf{p} = [1, z^{-1}, -z^{-1}, 1, -1]$. Applying Algorithm 2, we can obtain a 5×5 extension matrix P_e. The coefficient support of P_e satisfies $\text{coeffsupp}([\mathsf{P}_e]_{:,j}) \subseteq \text{coeffsupp}([\mathsf{p}_{\mathsf{a}_0}]_j)$ for $j = 1, \ldots, 5$. From P_e, we can derive high-pass filters $\mathsf{a}_1, \ldots, \mathsf{a}_4$ via (4.7) as follows:

$$\begin{aligned} \mathsf{a}_1(z) &= b_1(z) + b_1(z^{-1}); & \mathsf{a}_2(z) &= b_2(z) - b_2(z^{-1}); \\ \mathsf{a}_3(z) &= b_3(z) + z^3 b_3(z^{-1}); & \mathsf{a}_4(z) &= b_4(z) - z^3 b_4(z^{-1}); \end{aligned}$$

where

$$\begin{aligned}
b_1(z) \approx{}& -0.38736 + 0.26298i - (0.00307 + 0.00204i)z^6 - (0.00744 + 0.00660i)z^5\\
&- (0.00951 + 0.01459i)z^4 + (0.01372 - 0.02716i)z^3\\
&+ (0.06342 - 0.03967i)z^2 + (0.13657 - 0.04144i)z;\\
b_2(z) \approx{}& (0.00601 + 0.00672i)z^6 + (0.01353 + 0.02017i)z^5 + (0.01353 + 0.04034i)z^4\\
&- (0.01370 - 0.03024i)z^3\\
&- (0.07035 + 0.00011i)z^2 - (0.14727 + 0.03064i)z;\\
b_3(z) \approx{}& -(0.01063 + 0.01189i)z^6 - (0.02392 + 0.03567i)z^5 - (0.02392 + 0.07133i)z^4\\
&+ (0.05891 - 0.08433i)z^3 - (0.00043 - 0.20320i)z^2;\\
b_4(z) \approx{}& (0.00417 + 0.00467i)z^6 + (0.00939 + 0.01400i)z^5 + (0.00939 + 0.02800i)z^4\\
&- (0.02126 - 0.02655i)z^3 - (0.23403 + 0.31140i)z^2.
\end{aligned}$$

The coefficients of $b_1(z), \dots, b_4(z)$ are rounded from exact explicit solutions (too long to be presented here). We have $\mathsf{a}_1(z) = \mathsf{a}_1(z^{-1})$, $\mathsf{a}_2(z) = -\mathsf{a}_2(1/z)$, $\mathsf{a}_3(z) = \mathsf{a}_3(1/z)$, and $\mathsf{a}_4(z) = -z^3\mathsf{a}_4(z^{-1})$. Let ϕ be the 3-refinable function associated with the low-pass filter a_0. Let $\psi^1, \dots, \psi^4$ be the wavelet functions associated with the high-pass filters $\mathsf{a}_1, \dots, \mathsf{a}_4$ respectively by (1.4). Then $\phi(-\cdot) = \phi$, $\psi^1(-\cdot) = \psi^1$, $\psi^2(-\cdot) = -\psi^2$, $\psi^3(1-\cdot) = \psi^3$, and $\psi^4(1-\cdot) = -\psi^4$. See Figure 3 for the graphs of $\phi, \psi^1, \psi^2, \psi^3$ and ψ^4.



Figure 4.3: The graphs of $\phi, \psi^1, \psi^2, \psi^3, \psi^4$ (left to right, top to bottom). Real part:solid line. Imaginary part:dashed line.

Example 4. Consider dilation factor $\mathsf{d} = 3$. Let $m = 4, n = 2$, and the low-pass filter $a_0^{4,2}$ as in Example 2 with its symbol a_0 for the complex pseudo spline ϕ of order $(4,2)$ is given by

$$\mathsf{a}_0(z) = \left(\frac{\frac{1}{z} + 1 + z}{3}\right)^4 \left[\left(-\frac{4}{3} - \frac{2\sqrt{5}}{3}i\right)\frac{1}{z} + \left(\frac{11}{3} + \frac{4\sqrt{5}}{3}i\right) + \left(-\frac{4}{3} - \frac{2\sqrt{5}}{3}i\right)z\right].$$

By Theorem 8, the shifts of ϕ are linearly independent. Hence, there exist compactly supported dual refinable functions in $\widetilde{\phi} \in L_2(\mathbb{R})$ for ϕ, i.e., $\widehat{\widetilde{\phi}}(\mathrm{d}\xi) = \widetilde{a}_0(\xi)\widehat{\widetilde{\phi}}(\xi)$ for some low-pass filter $\widetilde{a}_0$ and $\langle \phi, \widetilde{\phi}(\cdot - k)\rangle = \delta_k$, $k \in \mathbb{Z}$. Here, we provide a low-pass filter $\widetilde{\mathsf{a}}_0(z)$ for $\widetilde{\phi}$ as follows:

$$\widetilde{\mathsf{a}}_0(z) = \left(\frac{\frac{1}{z} + 1 + z}{3}\right)^8 \left(b(z) + b(\frac{1}{z})\right),$$

where

$$\begin{aligned} b(z) = {} & \frac{329387}{2754} + \frac{209689}{1377} i\sqrt{5} - \left(\frac{102661}{816} + \frac{5464379}{22032} i\sqrt{5}\right) z \\ & - \left(\frac{177727}{2754} - \frac{551620}{4131} i\sqrt{5}\right) z^2 + \left(\frac{2967467}{22032} - \frac{1034833}{22032} i\sqrt{5}\right) z^3 \\ & + \left(-\frac{375253}{4131} + \frac{158555}{15147} i\sqrt{5}\right) z^4 + \left(\frac{24620753}{727056} - \frac{29059}{22032} i\sqrt{5}\right) z^5 \\ & - \frac{24103}{3366} z^6 + \left(\frac{11}{16} + \frac{21391}{727056} i\sqrt{5}\right) z^7. \end{aligned}$$

See Figure 4 for the graph of the 3-refinable function $\widetilde{\phi}$ associated with the low-pass filter $\widetilde{\mathsf{a}}_0(z)$. Note that $\widetilde{\phi} = \widetilde{\phi}(-\cdot)$.



Figure 4.4: The graph of $\widetilde{\phi}$. Real part:left. Imaginary part:right.

5. Conclusions

References

[1] C. K. Chui and J. A. Lian, Construction of compactly supported symmetric and antisymmetric orthonormal wavelets with scale = 3, *Appl. Comput. Harmon. Anal.*, **2** (1995), 21–51.

[2] L. H. Cui, Some properties and construction of multiwavelets related to different symmetric centers, *Math. Comput. Simul.*, **70** (2005), 69–89.

[3] I. Daubechies, Ten Lectures on Wavelets, SIAM, CBMS Series, 1992.

[4] I. Daubechies, B. Han, A. Ron, and Z. Shen, Framelets: MRA-based constructions of wavelet frames, *appl. comput. Harmon. Anal.*, **14** (2003), 1–46.

[5] B. Han, Symmetric orthonormal scaling functions and wavelets with dilation factor 4, *Adv. Comput. Math.*, **8** (1998), 221–247.

[6] B. Han, Matrix extension with symmetry and applications to symmetric orthonormal complex *M*-wavelets, *J. Fourier Anal. Appl.*, to appear.

[7] B. Han and H. Ji, Compactly supported orthonormal complex wavelets with dilation 4 and symmetry, *Appl. Comput. Harmon. Anal.*, **26** (2009), 422–431.

[8] W. Lawton, S. L. Lee, and Z. Shen, An algorithm for matrix extension and wavelet construction, *Math. Comp.*, **65** (1996), 723–737.

[9] A. Petukhov, Construction of symmetric orthogonal bases of wavelets and tight wavelet frames with integer dilation factor. *Appl. Comput. Harmon. Anal.*, **17** (2004), 198–210.

[10] A. Ron and Z. Shen, Affine systems in $L_2(\mathbb{R}^d)$: the analysis of the analysis operator. *J. Funct. Anal.* **148** (1997), 408–447.

[11] Z. Shen, Refinable function vectors, *SIAM J. Math. Anal.*, **29** (1998), 235–250.

[12] P. P. Vaidyanathan, Multirate Systems and Filter Banks, Prentice Hall, New Jersey, 1992.